

2018-07

Separate neural representations of prediction error valence and surprise: Evidence from an fMRI meta-analysis.

Fouragnan, Elsa

<http://hdl.handle.net/10026.1/11506>

10.1002/hbm.24047

Human Brain Mapping

Wiley

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

HUMAN BRAIN MAPPING**Separate neural representations of prediction error valence
and surprise: evidence from an fMRI meta-analysis**

Journal:	<i>Human Brain Mapping</i>
Manuscript ID	HBM-17-1205.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Fouragnan, Elsa; University of Oxford, Experimental Psychology Retzler, Chris; University of Huddersfield, Psychology Philiastides, Marios G; University of Glasgow,
Keywords:	meta-analysis, learning, decision making, reward, prediction error

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

**Title: Separate neural representations of prediction error valence and surprise:
evidence from an fMRI meta-analysis**

Short title: Separate neural correlates of prediction error valence and surprise

Authors: Elsa Fouragnan^{1,2}, Chris Retzler^{1,3} and Marios G. Philiastides¹

Affiliations: ¹Institute of Neuroscience & Psychology, University of Glasgow, Glasgow, UK.
²Department of Experimental Psychology, University of Oxford, Oxford, UK, ³Department of
Behavioural & Social Sciences, University of Huddersfield, Huddersfield, UK.

Acknowledgements: Funding for this work was provided by the Biotechnology and
Biological Sciences Research Council (BBSRC; grants BB/J015393/1–2 to M.G.P.) and the
Economic and Social Research Council (ESRC; grant ES/L012995/1 to M.G.P.). We also
thank Dr. Andrea Pisauero, Dr. Miriam Klein-Flugge and Dr. Jacqueline Scholl for helpful
comments on the manuscript.

Abstract

Learning occurs when an outcome differs from expectations, generating a reward prediction error signal (RPE). The RPE signal has been hypothesized to simultaneously embody the valence of an outcome (better or worse than expected) and its surprise (how far from expectations). Nonetheless, growing evidence suggests that separate representations of the two RPE components exist in the human brain. Meta-analyses provide an opportunity to test this hypothesis and directly probe the extent to which the valence and surprise of the error signal are encoded in *separate* or *overlapping* networks. We carried out several meta-analyses on a large set of fMRI studies investigating the neural basis of RPE, locked at decision outcome. We identified two valence learning systems by pooling studies searching for differential neural activity in response to *categorical* positive-vs-negative outcomes. The first valence network (negative > positive) involved areas regulating alertness and switching behaviors such as the midcingulate cortex, the thalamus and the dorsolateral prefrontal cortex whereas the second valence network (positive > negative) encompassed regions of the human reward circuitry such as the ventral striatum and the ventromedial prefrontal cortex. We also found evidence of a largely distinct surprise-encoding network including the anterior cingulate cortex, anterior insula and dorsal striatum. Together with recent animal and electrophysiological evidence this meta-analysis points to a sequential and distributed encoding of different components of the RPE signal, with potentially distinct functional roles.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

39 **Introduction**

40

41 Effective decision-making depends upon accurate outcome representations associated with
42 potential choices. These representations can be defined through reinforcement learning (RL)
43 [Rescorla and Wagner, 1972; Sutton, 1998], a modelling framework that uses the reward
44 prediction error (RPE), the difference between actual and expected outcomes, as a learning
45 signal to update future outcome expectations. In this framework, RPE is a signed quantity
46 and learning is driven by two separate components of the RPE signal: its *valence* (i.e. the
47 sign of the RPE, representing whether an outcome is better [+] or worse [-] than expected)
48 and its *surprise* (i.e. the modulus of the RPE, representing the degree [high or low] of
49 deviation from expectations). Whereas the valence informs an agent whether to reinforce or
50 extinguish a certain behaviour [Fouragnan et al., 2015; Fouragnan et al., 2017; Frank et al.,
51 2004], the surprise component determines the extent to which the strength of association
52 between outcome and expectations needs to be adjusted [Collins and Frank, 2016; Niv et
53 al., 2015; den Ouden et al., 2012].

54

55 This modelling framework has received considerable attention in neuroscience since the
56 early 90's when animal neurophysiological studies identified dopaminergic neurons in the
57 midbrain, in particular in the ventral tegmental area (VTA), the substantia nigra pars
58 compacta (SNc) and reticulata (SNr), whose tonic response profile appears to
59 simultaneously capture both components of the RPE signal outlined above [Montague et al.,
60 1996; Schultz et al., 1993; Schultz et al., 1997]. Specifically, these neurons show
61 anticipatory increase and suppression of their tonic activity in response to positive and
62 negative RPE respectively. While the anticipatory increase is proportional to the magnitude
63 of positive RPE, the magnitude of negative RPE is encoded by the duration of the basal
64 tonic suppression.

65

1
2
3 66 This discovery was a breakthrough in the field of learning and decision making and has
4
5 67 continued to be influential in the field over the past two and half decades (see [Schultz,
6
7 68 2016a; Schultz, 2016b] for a review). As a result, this neurophysiological work has strongly
8
9 69 motivated human functional magnetic resonance imaging (fMRI) research to identify the
10
11 70 corresponding macroscopic Blood-Oxygen-Level-Dependent (BOLD) pattern of the signed
12
13 71 RPE. This pattern of activity was expected to be such that the strength of the BOLD would
14
15 72 proceed from high positive RPEs > low positive RPEs > low negative RPEs > high negative
16
17 73 RPEs. More specifically, studies have employed a model-based fMRI approach, whereby
18
19 74 different types of reinforcement-learning models are first fitted to subjects' behavior to yield
20
21 75 parametric predictors for signed RPE against which fMRI data are subsequently regressed
22
23 76 [Daw et al., 2011; Fouragnan et al., 2013; Gläscher et al., 2010; O'Doherty et al., 2004;
24
25 77 O'doherty et al., 2007; Queirazza et al., 2017].
26
27 78

28
29 79 These fMRI studies have employed different algorithms to derive the signed RPE, ranging
30
31 80 from the simple formulation of the temporal difference learning algorithm to incorporating
32
33 81 action learning, notably using the Q-learning and SARSA ('state, action, reward, state, and
34
35 82 action') algorithms [Schonberg et al., 2010; Seymour et al., 2007; Tanaka et al., 2006].
36
37 83 According to qualitative reviews of this previous findings [O'doherty et al., 2007] as well as
38
39 84 quantitative, coordinate-based meta-analyses of these studies, the regions correlating with
40
41 85 the different formulations of signed RPE have been found to be predominantly subcortical,
42
43 86 including the striatum and amygdala, with some cortical regions, such as the ventromedial
44
45 87 prefrontal cortex and the cingulate cortex also reported [Bartra et al., 2013; Garrison et al.,
46
47 88 2013; Liu et al., 2011]. Additionally, substantial effort has been undertaken to identify how
48
49 89 different types of outcomes (primary reward such as food, or secondary reward such as
50
51 90 monetary outcomes) can modulate signed RPE in the same regions and the extent to which
52
53 91 it can be considered a domain-general, common currency signal [Sescousse et al., 2013].
54
55 92

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

93 While using trial-by-trial estimates of signed RPE from reinforcement-learning models has
94 provided an enormously productive framework for understanding learning and decision-
95 making, a growing number of studies have also discussed the complementary role of
96 surprise, namely the unsigned RPE, which can also be estimated at the single-trial level.
97 These include, but are not limited to, the use of trial-by-trial estimates of the modulus of RPE
98 or Bayesian surprise according to Bayesian learning theory [Hayden et al., 2011; Iglesias et
99 al., 2013]. Additionally, human electroencephalography (EEG) studies, attempting to offer a
100 temporal account of the cortical dynamics associated with RPE processing, did not find a
101 systematic monotonic response profile consistent with a single RPE representation but
102 instead offered evidence suggestive of separate representations for valence and surprise at
103 the macroscopic level of responses recorded on the scalp. Specifically, multiple recent EEG
104 studies combining model-based RPE estimates with single-trial analysis of the EEG revealed
105 an early outcome stage reflecting a purely categorical valence signal and a later processing
106 stage reflecting separate representations for valence and surprise [Fouragnan et al., 2015;
107 Fouragnan et al., 2017; Philiastides et al., 2010b]. These later valence and surprise signals
108 appeared in spatially distinct but temporally overlapping EEG signatures.
109
110 These findings suggest that, in addition to the fully monotonic firing pattern of midbrain
111 neurons, there exist individual representations for valence and surprise, potentially
112 subserving different functional roles during reward-based learning (e.g. approach-avoidance
113 behavior and the speed of learning via varying degrees of attentional engagement,
114 respectively). Here, we conducted an fMRI meta-analysis to explore the possibility that there
115 exist separate neuronal representations encoding valence and surprise promoting reward
116 learning in humans. We discuss the findings of our work in the context of recent reports from
117 animal neurophysiology and human neuroimaging experiments that provide evidence
118 towards a distributed coding of the different facets of the RPE signal [Brischoux et al., 2009;
119 Fouragnan et al., 2015; Fouragnan et al., 2017; Matsumoto and Hikosaka, 2009].

120

121 **Materials and Methods**

122

123 **Literature search.** We selected fMRI studies using the Pubmed database
124 (<http://www.ncbi.nlm.nih.gov/pubmed>) with the following search keywords: "(fMRI OR
125 neuroimaging) AND (prediction error OR reward OR surprise)" along with three initial filters
126 preselecting studies in which participants were human adults of over 19 years of age and
127 excluding reviews. This initial selection resulted in 724 candidates for inclusion to which a
128 further twenty papers were added from existing in-house reference libraries. Note that
129 previous meta-analyses used the terms "prediction error" or "reward" but we are the first to
130 include "surprise" in our systematic search for relevant papers [Bartra et al., 2013; Garrison
131 et al., 2013; Sescousse et al., 2013].

132

133 Abstracts from the 788 candidate-papers identified were then evaluated for inclusion in the
134 corpus according to the following criteria. We required studies of healthy human adults,
135 reporting changes in BOLD as a function of three different components of RPE: the
136 categorical valence, surprise and signed RPE, including statistical comparisons either in the
137 form of binary contrasts or continuous parametric analyses. Because the main objective of
138 the present meta-analysis is to examine the neural coding of RPE processing at decision
139 outcome, we also imposed the restriction that fMRI analyses were time-locked to the
140 presentation of outcomes (feedback). We used studies involving outcomes consisting of
141 abstract points, monetary payoffs, consumable liquids and arousing pictures but excluded
142 papers in which outcomes consisted of social feedback. We also required that studies used
143 functional brain imaging and did not use pharmacological interventions and ensured that the
144 reported coordinates were either in Montreal Neurological Institute (MNI) or Talairach space.
145 Finally, we excluded papers in which results were derived from region of interest (ROI) since
146 our meta-analytic statistical methods assume that foci are randomly distributed in the whole
147 brain under the null hypothesis. After applying these constraints our meta-analysis

1
2
3 148 comprised 102 publications with a total of 2316 participants, 144 contrasts, and 991
4
5 149 activation foci. The number of participants per study ranged from 8 to 66 (median = 24,
6
7 150 interquartile range [IQR] = 7).

8
9 151
10
11 152 **Study categorization.** The goal of this meta-analysis was to separately categorize studies
12
13 153 along the three components of RPE, locked at time of outcome, in order to: 1) identify the
14
15 154 extent to which there exist distinct neural representations for valence and surprise and 2)
16
17 155 identify whether the neural correlates of the signed RPE simply intersect those of valence
18
19 156 and surprise (possibly due to colinearities across these components) or appear as unique
20
21 157 clusters of activation reflecting the true combined influence of the two measures.

22 158
23
24 159 To group the relevant papers according to the three main RPE components we used the
25
26 160 following definitions: 1) *valence* represents the sign of the RPE and as such it is positive
27
28 161 when an outcome is better than expected and negative when worse than expected, 2)
29
30 162 *surprise* represents the absolute degree of deviation from expectations and is treated as an
31
32 163 unsigned quantity and 3) *signed RPE* simultaneously reflects the influence of both valence
33
34 164 and surprise and appears as a fully signed parametric signal. According to these definitions,
35
36 165 we identified several fMRI statistical analyses conducted in the original studies that fall under
37
38 166 each of the three RPE components (Table 1). The main assumptions of these fMRI
39
40 167 analyses, with regard to the BOLD signal as a function of each RPE component, are
41
42 168 presented schematically in Figure 1.

43 169
44
45 170 [Figure 1]
46
47 171
48
49
50 172 For the valence components, the literature has looked at neural responses which vary
51
52 173 categorically along positive-negative axes, as represented in patterns A (i) and (ii) of Figure
53
54 174 1. We therefore extracted activations exhibiting a relative BOLD signal increase for negative
55
56 175 relative to positive outcomes (NEG > POS: pattern A (i)) and greater BOLD for positive

relative to negative outcomes (POS > NEG: pattern A (ii)), respectively. We considered six types of fMRI statistical comparisons which reported coordinate results from either: (1) a contrast associated with negative > positive outcomes, (2) a contrast associated with negative > no outcomes, (3) a negative correlation with a trial-by-trial regressor modulated by [+1] for positive outcomes and [-1] for negative outcomes, (4) the positive correlation with the regressor described in (3), (5) a contrast associated with positive > negative outcomes and (6) a contrast associated with positive > no outcomes. We grouped results from contrasts 1-3 (i.e. NEG > POS) and contrasts 3-6 (i.e. POS > NEG) to capture regions yielding greater BOLD activity for negative relative to positive outcomes and a greater activity for positive relative to negative outcomes respectively (Table 1).

While the fMRI literature on RPE processing has produced a large amount of theoretical and empirical evidence for the valence and the signed RPE components, comparatively little has been done to directly investigate surprise as a separate component. Fewer studies have used fMRI regressors that were parametrically modulated by trial-to-trial changes in surprise using the unsigned RPE [Fouragnan et al., 2017; Hayden et al., 2011; Iglesias et al., 2013]. These studies used the terms "surprise", "unsigned RPE", or outcome "saliency" to refer to the mathematical modulus of RPE from computational learning models. In addition to these papers, our literature search has revealed a number of other measures (see below), which are highly correlated with outcome surprise, as defined by learning theory. We therefore used these measures as proxies of surprise to gain insights into the spatial extent of the relevant neural responses and the degree to which they overlap with those associated with valence.

Specifically, a recent line of research has investigated the neural basis of "Bayesian surprise" or "volatility", computed as the direct modulus of Bayesian predictive error [Ide et al., 2013; Iglesias et al., 2013; Mathys et al., 2014; O'Reilly et al., 2013] which correspond to the absolute difference between categorical outcomes and the probabilistic expectation of

1
2
3 204 these outcomes, estimated using Bayesian inference. In the framework of Bayesian learning,
4
5 205 the absolute Bayesian RPE plays an important role in learning from rapid changes in
6
7 206 behavioral exploration [Courville et al., 2006]. Finally, other studies used the term
8
9 207 “associability” which is a parameter in the Pearce-Hall model [Hall and Pearce, 1979; Pearce
10
11 208 and Hall, 1980] defined as the degree of divergence between an actual outcome and the
12
13 209 original expectation (e.g., the associative strength between a choice and an outcome). We
14
15 210 note however, that in the RL framework, associability can also refer to the learning rate. It is
16
17 211 clear from these reports that there is a lack of consistent terminology to refer to unsigned
18
19 212 RPE, which emphasizes the need for a more unified framework for studying RPE
20
21 213 processing.
22
23 214
24
25 215 To test for consistencies in the neuronal responses across these different reports, and
26
27 216 provide initial support for a unified representation of surprise, we grouped fMRI analyses
28
29 217 which reported outcome-locked activations resulting from: (1) a positive correlation with a
30
31 218 trial-by-trial regressor of the modulus (unsigned) RPE resulting from RL models across both
32
33 219 positive and negative outcomes ("surprise" or "unsigned RPE"), (2) a positive correlation
34
35 220 with a trial-by-trial regressor of the unsigned RPE resulting from Bayesian modelling
36
37 221 ("Bayesian Surprise" or "volatility"), (3) a positive correlation with a trial-by-trial regressor of
38
39 222 the free parameter of the Pearce-Hall model ("associability" term), (4) a contrast associated
40
41 223 with (high positive outcomes and high negative outcomes) > (low positive outcomes and low
42
43 224 negative outcomes OR no outcomes), (5) a positive correlation with a parametric regressor of
44
45 225 surprising positive RPE alone and (6) a positive correlation with a parametric regressor of
46
47 226 surprising negative RPE alone (Table 1). Figure 1 illustrates the hypothesized pattern of
48
49 227 BOLD signal predicted by these contrasts (pattern B), exhibiting a V shaped response profile
50
51 228 that is maximal for both highly surprising negative and positive RPEs. Despite possible
52
53 229 subtle differences in the definition of these measures we expected that only foci consistently
54
55 230 correlating with deviations from reward expectations would be revealed in this analysis.
56
57 231
58
59
60

One reason the surprise component has not been looked at closely in isolation is because the literature has focused primarily on signed RPE representations instead. This approach was motivated by neurophysiology experiments showing monotonic responses as a function of both valence and surprise and by a theoretical framework suggesting that learning is driven by a single signed RPE representation. To identify the spatial extent of these representations we also looked at fMRI data reporting positive correlations with signed RPE (negative correlation were discarded). Specifically, we combined four types of fMRI analyses, which estimated trial-by-trial signed RPE from different computational models. We used fMRI reports from (1) model-free and (2) model-based RL methods. Model-free methods include Markov Chain Monte Carlo and temporal difference methods [Samson et al., 2010; Seymour et al., 2007]. Model-based methods include dynamic programming and certainty equivalent methods [Daw et al., 2005; Doya et al., 2002]. More on these algorithms can be found in the review by [Kaelbling et al., 1996]. We also included continuous parametric analyses using trial-by-trial signed RPE from (3) Bayesian RL framework described above [Iglesias et al., 2013; Mathys et al., 2014; den Ouden et al., 2012]. Finally, our analysis for signed RPE also contained one type of parametric analysis that employed fixed RPE values (not estimated from RL models) ranked on a scale such that (4) high positive RPEs > low positive RPEs > low negative RPEs > high negative RPEs (Table 1). Figure 1 illustrates the hypothesized pattern of BOLD signal predicted by these contrasts (pattern C) and it is assumed to increase linearly as a function of signed RPE.

Crucially, we note that an issue requiring closer scrutiny pertains to the difficulty in disambiguating the signed RPE pattern of activity from those associated with valence and surprise. Specifically, pattern C (signed RPE) is generally highly correlated with pattern A (ii), (POS > NEG valence) and in studies in which only positive RPEs are considered, pattern C (signed RPE) and pattern B (surprise) are perfectly correlated. Nonetheless, comparing clusters of activations across the three RPE components could potentially reveal whether or not there exist unique clusters of activations associated with signed RPE.

1
2
3 260
4
5 261 [Table 1]
6
7 262
8
9 263
10
11 264
12
13 265
14
15 266
16
17 267
18
19 268
20
21 269
22
23 270
24
25 271
26
27 272
28
29 273
30
31 274
32
33 275
34
35 276
36
37 277
38
39 278
40
41 279
42
43 280
44
45 281
46
47 282
48
49 283
50
51 284
52
53 285
54
55 286
56
57
58
59
60

2.1. Activation Likelihood Estimation (ALE) analysis

We conducted the meta-analysis using the GingerALE software (version 2.3.6) [Eickhoff et al., 2009] that employs a revised (and rectified [Eickhoff et al., 2017]) version of the activation likelihood estimation (ALE) algorithm [Laird et al., 2005; Turkeltaub et al., 2002], which identifies common areas of activation across studies. This method performs coordinate based meta-analysis which considers each reported foci as a 3D Gaussian probability distribution, centred at the coordinates provided by each study reflecting the spatial uncertainty associated with each reported set of coordinates. Note that each contrast provided to the ALE algorithm is treated as a separate experiment. The probabilities distributions are then combined to create a modelled activation map, namely an ALE map for that contrast. Studies are weighted according to the number of subjects they contain by adjusting the full width at half maximum of the Gaussian distributions. The convergence of results across the whole brain is obtained by computing the union of all resulting voxel-wise ALE scores. To distinguish meaningful convergence from random noise, statistics are computed by comparing ALE scores with an empirical null-distribution representing a random spatial association between studies. To infer true convergence, a random-effect inference is applied to capitalize on the differences between studies rather than between foci within a particular study. The null-hypothesis is modelled by randomly sampling voxels from each of the ALE maps from which the union is obtained. The ALE maps are assessed against the null distribution using a cluster level threshold of specific p-values. Contrast analyses between categories of the entire dataset are determined by ALE subtraction method, including a correction for differences in sample size between the categories.

Here, we manually extracted all coordinates from the studies shown in Table 1 and entered them into separate files for each of the three RPE components in preparation for the ALE analyses. Any studies that provided coordinates in Talairach space were converted into MNI space by the Matlab (MathWorks, Natick, Massachusetts) function *tal2mni* in the fieldtrip toolbox [Oostenveld et al., 2011]. We conducted ALE analyses for each of the three components of RPE individually. Along the valence component, we looked at both patterns A (i) and A (ii) in Figure 1 (i.e. to identify activations for negative > positive RPE and vice versa, respectively). Accordingly, we ran separate ALE analyses for each of the two patterns. In addition, we performed two conjunction analyses – one between the valence and surprise components to investigate our hypothesis of largely separate neural representations and another between all three RPE components to identify regions that simultaneously encode these representations. Subsequently, we also performed all possible pairwise contrast analyses between the three patterns (A, B and C), using the individual maps associated with each pattern.

A total of 402 foci from 66 contrasts were used with 262 foci from 31 contrasts for Pattern A (i) revealing BOLD patterns greater for negative than positive outcomes and 205 from 35 contrasts for Pattern A (ii) (e.g. the opposite contrast). For the surprise (Pattern B) and signed RPE (Pattern C) analyses, we applied individual ALE analyses, with 284 foci from 40 contrasts for surprise and 240 foci from 38 contrasts for signed RPE. Overall, the number of contrasts used for each separate outcome component was large enough (> 30) to allow sufficient power for the required statistical tests [Eickhoff and Etkin, 2016]. Finally, we transformed the resulting ALE maps from the Colins MNI individual brain space (Colin27_T1_seg_MNI) to the MNI normalized brain space (MNI ICBM152 template) by applying an affine transformation using the FSL *flirt* program [Jenkinson et al., 2002], prior to overlaying onto the canonical MNI template for visualization.

312

313 3. Results

1
2
3 314
4
5 315 All coordinates used for the following ALE analyses were collated from fMRI studies in which
6
7 316 the components of RPE have been regressed onto BOLD activity time-locked to outcome
8
9 317 presentation. We report ALE maps with clusters surviving the False Discovery Rate (FDR)
10
11 318 yielding two p-value thresholds. The most conservative FDR correction yields a p-value with
12
13 319 no assumptions about how the data is correlated (FRN), and the least conservative FDR
14
15 320 correction assumes independence or positive dependence (FID) with $p < 0.05$ and a
16
17 321 minimum volume clustering value of 50 mm^3 . Note that, using a cluster-level family-wise
18
19 322 error (FWE) correction implemented with a cluster-extent threshold of $p < 0.05$ and a cluster-
20
21 323 forming threshold of $p < 0.001$ revealed virtually identical results (compared with FRN)
22
23 324 [Eickhoff et al., 2017] as per previous reports [R Garrison et al., 2017]. For all tables
24
25 325 presenting ALE cluster results, the size of each cluster is provided in mm^3 along with the
26
27 326 associated MNI coordinates and maximum ALE score. The ALE score indicates the relative
28
29 327 effect size for each peak voxel within each ALE analysis.

30
31 328
32
33 329 **3.1. Outcome Valence**

34
35 330
36
37 331 The first two ALE analyses were conducted to identify regions in which BOLD signals
38
39 332 correlate with outcome valence. Specifically, we looked at activations that yielded greater
40
41 333 BOLD for negative relative to positive outcomes (NEG > POS; pattern A (i) in Figure 1) and
42
43 334 greater BOLD for positive relative to negative outcomes (POS > NEG; pattern A (ii) in Figure
44
45 335 1), respectively. Accordingly, we considered all fMRI studies, which assumed BOLD
46
47 336 responses varying categorically along a positive-negative axis for outcome valence.

48
49 337
50
51 338 The findings of the two valence ALE analyses are shown in Figure 2. The resulting maps
52
53 339 revealed a highly distributed network of brain activations encompassing several cortical
54
55 340 regions and sub-cortical structures. More precisely, NEG > POS valence clusters were found

in a network encompassing the anterior and dorsal part of the mid-cingulate cortex (aMCC and dMCC) including the pre supplementary motor area (pre-SMA), the bilateral anterior and middle insular cortex (aINS, mINS), the bilateral dorsolateral prefrontal cortex (dlPFC), the bilateral thalamus, right amygdala, left inferior parietal lobule (IPL) and the habenula.

POS > NEG valence clusters were found in the bilateral ventral striatum (vSTR), the ventromedial prefrontal cortex (vmPFC), the posterior part of the cingulate cortex (PCC), as well as the ventrolateral orbitofrontal cortex (vlOFC). At a lower threshold (uncorrected p-value of 0.001), we also found the midbrain as part of this network, encompassing the VTA, which is commonly associated with the delivery of reward [D'Ardenne et al., 2008]. Table 2 contains the complete list of regions, coordinates, and statistics of these two ALE analyses.

[Figure 2], [Table 2]

3.2. Surprise

FMRI investigations of RPE have focused primarily on the valence components while neglecting potential contributions from possible separate representations along the surprise component, defined as the degree by which outcomes deviate from expectations and mathematically expressed as the modulus of RPE. A major goal of this work was to explore the possibility that there exist largely separate neuronal representations encoding surprise. To this end, we conducted a new ALE analysis in which the few empirical fMRI studies making use of the surprise from RL models were combined with other fMRI measures correlated with the surprise as defined by RL models (Table 1).

Figure 3 shows the areas in which BOLD signal correlated with surprise. We found evidence for activations in a distributed network encompassing the aMCC, dMCC, the pre-SMA the bilateral dorsal striatum (dSTR), the bilateral aINS, the MTG and the midbrain. Crucially, this

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

activation map shows that the neural network associated with surprise is largely distinct from that of valence. This finding provides initial support for the notion that these two RPE components are encoded in separate brain areas and, as such, they might be contributing individually to promote learning. The full results of the surprise ALE analysis are also summarized in Table 3.

[Figure 3], [Table 3]

3.3. Valence and surprise conjunction and contrast analyses

The activation maps for valence (NEG > POS and POS > NEG) and surprise ALE analyses conducted above revealed little overlap between the spatial representations of these two RPE components. To formally quantify the degree of overlap between the valence and surprise networks, we next ran a conjunction analysis between the two components. The statistical map resulting from this conjunction analysis and the two separate statistical maps of valence and surprise (as already reported in Figures 2 and 3) are overlaid in Figure 4.

[Figure 4], [Table 4]

Contrast analyses were conducted for each possible pairing between any dimensions of valence (POS > NEG [positive]; NEG > POS [negative] and POS + NEG [all valence]) and surprise. These analyses allowed us to identify the areas that were unique and specific to each individual outcome and RPE-related component. The positive valence (pattern A (ii)) minus surprise (pattern B) contrast revealed two main clusters in the vSTR and vmPFC whereas the reverse contrast revealed a network of clusters including preSMA, aINS, and MTG. Contrasting negative valence (pattern A (i)) and surprise also exposed separate networks of areas for each subtraction. Specifically, this contrast revealed a network encompassing the thalamus, the habenula, the right mINS and the dMCC, whereas the

reverse contrast showed clusters in the dorsal portion of the STR and the dlPFC. The statistical maps resulting from these contrast analyses are presented in Figure 5.

[Figure 5], [Table 5]

3.4. Signed RPE

A major goal of this work was to investigate the spatial profile of the signed RPE component and to scrutinise more closely the extent to which it overlaps with the separate representations identified for valence (NEG > POS and POS > NEG) and surprise. The fMRI-RPE literature has focused on this component largely due to neurophysiological evidence suggesting that RPE-like learning is driven by a single, theoretically unified representation of both POS > NEG valence and surprise (Table 1).

Results from this ALE analysis revealed very few unique activations for signed RPE compared to valence and surprise. Instead, brain areas identified in this analysis overlapped mostly with areas appearing in the POS > NEG valence component and, to a lesser extent, surprise (Figure 6). Specifically, a large overlap between signed RPE and the POS > NEG valence component was found in the STR and a smaller one in the vmPFC. Similarly, areas appearing in the signed RPE analysis that overlapped with the surprise component were also found, albeit only in small clusters comprising the aMCC and dorsal STR. Taken together, these findings emphasize the potential collinearities between the BOLD predictors used to identify neural representations associated with the three RPE components and highlight the need for developing a methodology for properly disentangling their individual contributions.

[Figure 6], [Table 6]

1
2
3 425 **3.5. Putting it all together**

4 426
5
6
7 427 Subsequently, to formally test for the overlap between all three RPE components and
8
9 428 identify potential regions integrating valence and surprise either into a signed RPE
10
11 429 representation or a linear superposition of the two signals [Fouragnan et al., 2017], we
12
13 430 performed a conjunction analysis between the valence (pattern A), the surprise (pattern B)
14
15 431 and signed RPE (pattern C) signals. We summarize our conjunction results in Figure 7,
16
17 432 which revealed a major overlap between all activations associated with signed RPE and
18
19 433 each of the other two RPE representations in the central part of the STR. Thus, one
20
21 434 possibility is that the STR meets the requirement that a full monotonic representation of the
22
23 435 error signal also simultaneously encodes valence and surprise, as per our last ALE analysis.

24 436
25
26 437 [Figure 7]
27
28 438

29
30 439 Another possibility is that the overlap between all components of outcomes in the STR is
31
32 440 arising, at least in part, due to collinearities across the different outcome representations,
33
34 441 particularly between the positive categorical nature of outcome valence (pattern A (ii)) and
35
36 442 the signed RPE. To formally test this hypothesis, we performed a new series of contrast
37
38 443 analyses between signed RPE and all dimensions of categorical valence and surprise.
39
40 444 Particularly, we performed contrast analyses between patterns C-A(i), C-A(ii), C-A and C-B
41
42 445 (and vice versa). The results are summarized in Figure 8. Particularly, we did not find any
43
44 446 area unique to signed RPE when looking at each of the individual comparisons of signed
45
46 447 RPE with the other three patterns. In fact, when comparing signed RPE to positive valence
47
48 448 (pattern A (ii)), no clusters were found to be significantly different than those found with the
49
50 449 categorical outcome valence (POS > NEG). Conversely, the STR was found for all the other
51
52 450 signed RPE comparisons (signed RPE > negative; signed RPE > surprise). Finally, the
53
54 451 unique network related to negative valence (pattern A (i)) was found in the dMCC, thalamus

and mINS, the unique cluster related to positive valence was found in the vmPFC and the unique network related to surprise was found in the aMCC, preSMA and the aINS.

[Figure 8], [table 7]

Discussion

In this fMRI meta-analysis work, we demonstrated that reward learning in humans involves separate neuronal signatures of RPE, comprising distinct representations for valence and surprise. Together with recent neurophysiological and EEG evidence (including studies using simultaneous EEG and fMRI), these findings point to a potentially sequential and distributed encoding of different RPE components with potentially functionally distinct roles.

Valence networks

The ALE analyses related to valence revealed two distributed set of activations correlating with both pattern A (i) and (ii) in Figure 1. Foci for which the BOLD signal was greater for negative than positive outcomes showed significant clustering in a large network of areas including the thalamus, the aMCC and dMCC, the aINS, mINS and the dIPFC. Conversely, foci for which the BOLD signal was greater for positive than negative outcomes showed significant clusters in a separate network including vmPFC, vSTR, PCC, and vIOFC. These findings clearly suggest the presence of multiple systems responding to the categorical nature of valence which supports the notion that separate valuation systems shape learning in the human brain [Fiorillo, 2013; Fouragnan et al., 2013], although their functional role remain debated. More specifically, the debate focuses on the number and exact nature of the neural systems assigning value to decision outcomes and driving behaviors that are evolutionarily appropriate in response to changes in the environment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

480 A first theory describes two distinct valence systems invoking two orthogonal axes of
481 decision-making: alertness (involving the implementation of action) and learning (including
482 the updates of value expectations for future avoidance and approach behaviors). In this
483 framework, the first system is thought to monitor on-going activity and interrupt it when
484 needed to trigger switching behaviors (e.g. following negative RPEs). In contrast, the second
485 system uses both negative and positive RPE values for decreasing or increasing internal
486 value representations associated with decisions to ultimately drive avoidance and approach
487 learning, respectively [Boureau and Dayan, 2011; Cools et al., 2011; Elliot, 2006; Fiorillo,
488 2013; Fouragnan et al., 2015; Gray and McNaughton, 2003; Guitart-Masip et al., 2012].

489
490 A second (not mutually exclusive) proposition supports the idea that there are at least two
491 separate systems responsible for aversive and appetitive reinforcements such that
492 punishments and rewards are encoded separately (i.e. a punishment space and a reward
493 space [Morrens, 2014]). This proposition was developed on the basis of neurophysiological
494 evidence showing that different types of neurons exhibit differential activity in response to
495 punishing vs. non-punishing outcomes and rewarding vs. non-rewarding outcomes,
496 respectively [Fiorillo et al., 2003; Fiorillo, 2013; Schultz et al., 1992; Schultz, 1998]. In this
497 second theory, the punishment space is responsible for avoidance behaviors as well as
498 avoidance learning and the reward space is responsible for approach behaviors and
499 approach learning.

500
501 It is noteworthy that our meta-analysis on itself cannot directly distinguish between the two
502 theories because the results do not reveal whether the relevant activations respond
503 exclusively to either positive or negative outcomes or are modulated by both outcomes in
504 opposite directions. This distinction is critical because the former response profile would
505 suggest the presence of separate approach and avoidance systems that might not
506 necessarily be linked to the learning processes as such, while the latter might point to both
507 up- and down-regulation of activity consistent with learning and updating of reward

expectations. Nonetheless, the meta-analysis results suggest that two main networks process valence. The network encompassing aINS, aMCC, thalamus and dlPFC could regulate on-going activity and alertness or could represent the punishment space in accordance to the first and the second theories respectively. Conversely, the network of regions encompassing the vmPFC, vSTR, PCC and vIOFC could represent the learning system depicted in the first theory or could represent the reward space depicted in the second theory. Further research is required to tease apart the roles of these systems, especially by investigating their precise response profiles in the appetitive (where rewarding and non-rewarding outcomes are manipulated) and in a true aversive (where punishing and non-punishing outcomes are manipulated) domains, respectively.

Surprise network

Emerging evidence indicates that the brain encodes the unsigned RPE signal (surprise), which alerts the organism of relative deviations from expectations, regardless of the outcome value. However, to date, only few papers have modelled surprise as such to search for independent neural representations, with the exception of recent neurophysiological developments [Brischoux et al., 2009; Matsumoto and Hikosaka, 2009], recent EEG work [Philiastides et al., 2010b; Yeung and Sanfey, 2004] and an increasing number of fMRI studies [Fouragnan et al., 2017; Gläscher et al., 2010; Li and Daw, 2011; Metereau and Dreher, 2013]. Nevertheless, other fMRI studies used variables highly correlated with surprise that can be employed as proxies [Behrens et al., 2007; Iglesias et al., 2013; Nassar et al., 2012; den Ouden et al., 2012; Yu and Dayan, 2005]. These studies share the assumption that the corresponding BOLD response profile is maximal for high positive and high negative RPE and minimal for no RPE, resembling a V-shape, as illustrated with Pattern B in Figure 1. By combining these fMRI results into a single ALE-analysis, we expose for the first time the network associated with surprise while stressing the need for a common lexicon for this learning component to guide subsequent research in the field.

1
2
3 536
4
5 537 The surprise ALE-analysis revealed a large network including cortical and sub-cortical areas
6
7 538 such as aMCC, bilateral aINS, dSTR and midbrain, that differed majoritarily from those of
8
9 539 valence processing although small overlaps were found between the two components at the
10
11 540 junction of ventral and dorsal STR, in left aINS and aMCC. Importantly, the role of surprise is
12
13 541 still a subject of debate. Some studies propose that this network encodes the saliency of an
14
15 542 outcome or how much a stimulus stands out from others [Litt et al., 2011; Zink et al., 2004].
16
17 543 As such, the surprise system could be considered as a key attentional mechanism that
18
19 544 enables an organism to focus its limited perceptual and cognitive resources on the most
20
21 545 pertinent subset of the available sensory data, similarly to the attentional mechanism used to
22
23 546 guide decisions in the case of salient stimuli [Kahnt and Tobler, 2013]. Consistent with a role
24
25 547 in attention regulation, representations of such signal have been found in lower-level visual
26
27 548 areas [Serences, 2008], lateral intraparietal cortex [Huettel et al., 2006; Kahnt and Tobler,
28
29 549 2013] and areas involved in visual and motor preparation such as the supplementary motor
30
31 550 area [Wunderlich et al., 2009] or the supplementary eye field [Middlebrooks and Sommer,
32
33 551 2012; So and Stuphorn, 2012].
34
35 552
36
37 553 In contrast, it has also been suggested that a surprise system can independently monitor
38
39 554 unexpected information and act as a learning signal that allows better predictions of
40
41 555 upcoming events, and help plan appropriate behavioral adjustments [Dayan and Balleine,
42
43 556 2002; Fouragnan et al., 2017; Kolling et al., 2012; Wittmann et al., 2016]. In particular, some
44
45 557 studies suggest that the aINS receives information related to surprise and direct modulation
46
47 558 from the dSTR providing crucial information for behavioral adjustment [Menon and Levitin,
48
49 559 2005]. Along these lines, the surprise signal also captures the essence of a learning signal
50
51 560 that the brain needs to compute to maintain a homeostatic state [Friston et al., 2006; Friston,
52
53 561 2009]. Practically, this means that the brain elaborates internal predictions about sensory
54
55 562 input and updates them according to surprise, a process that can be formulated as
56
57 563 generalized Bayesian filtering or predictive coding in the brain. Finally, still in the framework
58
59
60

of learning, some authors argue that surprise can also be considered as a signal predicting the level of risk associated with a future decision outcome, and thus reflect a risk RPE [Fiorillo et al., 2003; Preuschoff et al., 2008; Rudolf et al., 2012].

Neuromodulatory pathways encoding multicomponent RPE signals

Supporting the idea of separate neural systems for valence and surprise, recent electrophysiological work has revealed both signals existing in neighbouring groups of neurons. The first study of this kind observed the response of dopaminergic neurons in ventral and dorsal areas of the SNc and reported two categories of dopamine neurons [Matsumoto and Hikosaka, 2009]. Some dopamine neurons increase their phasic firing activity in response to valence while others responded only to the changes in unsigned RPE, regardless of the valence component. The latter population of neurons was located more dorsolaterally in the SNc, whilst the neurons encoding valence were located more ventromedially, including the VTA. Interestingly, the dorsolateral SNc projects mainly to the dorsal STR, whereas the ventral SNc and VTA project to the ventral STR, which matches the results of our last conjunction analysis (Figure 7). We found that the only region that encodes the full monotonic representation of the RPE as well as the separate valence and surprise components of RPE seems to be the central part of the STR as shown in Figure 7. This result aligns with the assumption that this region receives direct projections from the midbrain dopaminergic neurons encoding a fully monotonic signed RPE signal [Schultz et al., 1997]. Additionally, the meta-analysis also revealed that both the valence (POS > NEG) and surprise networks include activity in the midbrain, confirming this hypothesis.

It is important to note that identifying neural activity associated with valence and surprise signals is challenging because in many experimental paradigms both components are highly correlated. For example, when positive RPE are manipulated in isolation, valence (POS > NEG) strongly correlates with surprise. Additionally, whether positive or negative, an

1
2
3 592 unexpected outcome attracts more attention, leads to higher levels of emotional arousal and
4
5 593 involves higher levels of motor preparation compared to no RPE [Matsumoto and Hikosaka,
6
7 594 2009; Maunsell, 2004; Roesch and Olson, 2004]. Consequently, to disentangle these
8
9 595 signals, one needs to design tasks in which the level of valence and surprise can
10
11 596 independently be controlled and decoupled [Kahnt, 2017; Kahnt and Tobler, 2013] or
12
13 597 capitalize on the variability of physiologically-derived responses (i.e. endogenous variability)
14
15 598 associated with valence and surprise [Fouragnan et al., 2015; Fouragnan et al., 2017;
16
17 599 Pisauro et al., 2017].

18
19 600
20
21 601 It is important to note that since the problem of collinearity and functional specificity of some
22
23 602 brain regions is already present in single studies, it will inevitably be carried over to studies
24
25 603 performing conjunction meta-analyses. Virtually every experimental design engages a large
26
27 604 number of cognitive operations and, thereby, activates functional neural networks that may
28
29 605 be irrelevant to a particular regressor (psychological construct) of interest. For example in
30
31 606 our study, regions related to outcome value and surprise might share variance with outcome
32
33 607 confidence [Gherman and Philiastides, 2015; Gherman and Philiastides, 2017; Lebreton et
34
35 608 al., 2015; Philiastides et al., 2014]. Despite this general limitation and the difficulty of
36
37 609 interpreting conjunction results, aggregating results across a large number of experiments
38
39 610 allows one to expose convergence of findings across studies and increasing the
40
41 611 generalizability of the conclusions. In particular, this meta-analysis, capitalizing on both
42
43 612 individual maps of activations as well as contrasts between different outcome components,
44
45 613 points to a distributed encoding of valence and surprise, with potentially distinct functional
46
47 614 roles.

48 615
49
50 616 **Temporally specific components of RPE processing**

51
52 617
53
54 618 The presence of separate RPE-related neural systems raises the question of how these
55
56 619 systems unfold in time. Capitalizing on the high temporal resolution of EEG, three recent

studies using simultaneous EEG-fMRI have started to shed light on the spatiotemporal characterisation of the RPE components. First, these studies have revealed two temporally specific EEG components discriminating between positive and negative RPEs peaking around 220ms and 300ms respectively, largely consistent with the timing of the feedback-related negativity and feedback-related positivity ERP components [Cohen et al., 2007; Hajcak et al., 2006; Yeung and Sanfey, 2004]. Additionally, the studies also revealed a late unsigned RPE component which overlaps temporally with the late valence signal [Philiastides et al., 2010b] but appears in a largely separate and distributed neural network [Fouragnan et al., 2017].

Based on these previous studies and the current meta-analysis, we propose that the early and late EEG valence components might reflect the separate contributions of the two networks of areas found for the ALE-valence analyses. This proposal assumes that an early network processes mainly negative RPEs in order to initiate a fast alertness response in the presence of negative outcomes. Conversely, a later network – associated with the brain's reward circuitry – is modulated by both positive and negative RPEs, consistent with a role in approach/avoidance learning and value updating [Philiastides et al., 2010a]. We also propose that the surprise network unfolds near simultaneously with the late valence component and thus influences learning through largely distinct spatial representations of the two outcomes signals, which happen to form a composite signal in overlapping areas [Fouragnan et al., 2017].

641

642 **Full representation of a monotonic signed RPE signal**

643

To examine the spatial profile of a true monotonic signed RPE representation in the human brain, we pooled results from fMRI studies, which hypothesized that RPE-like learning is driven by a simultaneous representation of both categorical valence and surprise. These

1
2
3 647 fMRI studies are based on the influential assumption that BOLD signal increases
4
5 648 monotonically as a function of signed RPE, as illustrated in pattern C (Fig. 1), equivalent to
6
7 649 the teaching signal that is predicted in the Rescorla–Wagner model of RL [Rescorla and
8
9 650 Wagner, 1972]. Additionally, we combined the valence and surprise networks and
10
11 651 subsequently compared it with the signed RPE to test the requirement that the signed RPE
12
13 652 simultaneously encodes both components. This conjunction analysis revealed that the only
14
15 653 brain region that seems to encode a true monotonic signal is the STR in the basal ganglia,
16
17 654 which could explain why such a signal is not tractable with EEG recordings as highlighted
18
19 655 earlier. This result confirms the long standing view that the BOLD activity in STR mirrors the
20
21 656 dopaminergic signalling of the mesolimbic neurons [Delgado et al., 2000; Haber et al., 1995;
22
23 657 O'Doherty et al., 2004; Pagnoni et al., 2002] that fully encode the RL prediction error signal
24
25 658 of the Rescorla-Wagner rule [Ikemoto, 2007; Schultz et al., 1992].
26

27 659
28
29 660 Nonetheless, the ALE contrast analyses between valence (the positive correlation with
30
31 661 pattern A (ii)) and signed RPE revealed no significant activation, whereas the reverse
32
33 662 contrast revealed a denser cluster of activity in vmPFC for valence than signed RPE. Given
34
35 663 the evidence presented above that the signed RPE may only be encoded in the STR, we
36
37 664 suggest that this result may arise due to collinearities between valence and signed RPE or
38
39 665 surprise and signed RPE. More precisely, a parametric predictor for signed RPE would be
40
41 666 positively correlated with the contrast positive > negative outcomes whereas the signed RPE
42
43 667 and surprise would be perfectly correlated in the positive (appetitive) domain.
44

45 668
46 669 **Conclusion**
47

48 670
49
50 671 In conclusion, the current meta-analysis points to a framework whereby heterogeneous
51
52 672 signals are involved in RPE processing. The proposal of a temporally distinct and spatially
53
54 673 distributed representation of valence and surprise is open to debate and many questions
55
56 674 remain about how these signals interact and how they correspond to the computations made
57
58
59
60

1
2
3 675 in the brain. For example, it is currently unclear whether valence and surprise encoding
4
5 676 occur before the computation of the signed RPE, or whether these three computations are
6
7 677 performed in parallel. Nevertheless the taxonomy proposed is conceptually useful because it
8
9 678 breaks down the learning and valuation processes into testable components and organizes
10
11 679 the RPE literature in terms of the computations that are potentially involved. It will require
12
13 680 additional experiments to validate the current proposal and to better understand the
14
15 681 complexity of RPE processing.
16
17 682

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

Abler B, Walter H, Erk S, Kammerer H, Spitzer M (2006): Prediction error as a linear function of reward probability is coded in human nucleus accumbens. *NeuroImage* 31:790–795.

Allen M, Fardo F, Dietz MJ, Hillebrandt H, Friston KJ, Rees G, Roepstorff A (2016): Anterior insula coordinates hierarchical processing of tactile mismatch responses. *NeuroImage* 127:34–43.

Amado C, Hermann P, Kovács P, Grotheer M, Vidnyánszky Z, Kovács G (2016): The contribution of surprise to the prediction based modulation of fMRI responses. *Neuropsychologia* 84:105–112.

Amiez C, Sallet J, Procyk E, Petrides M (2012): Modulation of feedback related activity in the rostral anterior cingulate cortex during trial and error exploration. *NeuroImage* 63:1078–1090.

Aron AR, Shohamy D, Clark J, Myers C, Gluck MA, Poldrack RA (2004): Human Midbrain Sensitivity to Cognitive Feedback and Uncertainty During Classification Learning. *J Neurophysiol* 92:1144–1152.

Bartra O, McGuire JT, Kable JW (2013): The valuation system: a coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *NeuroImage* 76:412–427.

Behrens TEJ, Woolrich MW, Walton ME, Rushworth MFS (2007): Learning the value of information in an uncertain world. *Nat Neurosci* 10:1214–1221.

Bickel WK, Pitcock JA, Yi R, Angtuaco EJC (2009): Congruence of BOLD Response across Intertemporal Choice Conditions: Fictive and Real Money Gains and Losses. *J Neurosci* 29:8839–8846.

Boll S, Gamer M, Gluth S, Finsterbusch J, Büchel C (2013): Separate amygdala subregions signal surprise and predictiveness during associative fear learning in humans. *Eur J Neurosci* 37:758–767.

van den Bos W, Cohen MX, Kahnt T, Crone EA (2012): Striatum-medial prefrontal cortex connectivity predicts developmental changes in reinforcement learning. *Cereb Cortex N Y N 1991* 22:1247–1255.

Boureau Y-L, Dayan P (2011): Opponency Revisited: Competition and Cooperation Between Dopamine and Serotonin. *Neuropsychopharmacology* 36:74–97.

Brischoux F, Chakraborty S, Brierley DI, Ungless MA (2009): Phasic excitation of dopamine neurons in ventral VTA by noxious stimuli. *Proc Natl Acad Sci U S A* 106:4894–4899.

Browning M, Holmes EA, Murphy SE, Goodwin GM, Harmer CJ (2010): Lateral Prefrontal Cortex Mediates the Cognitive Modification of Attentional Bias. *Biol Psychiatry* 67:919–925.

de Bruijn E, de Lange F, Cramon D, Ullsperger M (2009): When Errors Are Rewarding. *J Neurosci Off J Soc Neurosci* 29:12183–6.

Canessa N, Crespi C, Motterlini M, Baud-Bovy G, Chierchia G, Pantaleo G, Tettamanti M, Cappa SF (2013): The Functional and Structural Neural Basis of Individual Differences in Loss Aversion. *J Neurosci* 33:14307–14317.

Chumbley JR, Burke CJ, Stephan KE, Friston KJ, Tobler PN, Fehr E (2014): Surprise beyond prediction error. *Hum Brain Mapp* 35:4805–4814.

- 1
- 2
- 3 727 Cohen MX, Elger CE, Ranganath C (2007): Reward Expectation Modulates Feedback-Related
- 4 728 Negativity and EEG Spectra. *NeuroImage* 35:968–978.
- 5 729 Cohen MX, Ranganath C (2007): Reinforcement Learning Signals Predict Future Decisions. *J*
- 6 730 *Neurosci* 27:371–378.
- 7 731 Collins AGE, Frank MJ (2016): Surprise! Dopamine signals mix action, value and error. *Nat*
- 8 732 *Neurosci* 19:3–5.
- 9 733 Cools R, Nakamura K, Daw ND (2011): Serotonin and Dopamine: Unifying Affective,
- 10 734 Activational, and Decision Functions. *Neuropsychopharmacology* 36:98–113.
- 11 735 Courville AC, Daw ND, Touretzky DS (2006): Bayesian theories of conditioning in a changing
- 12 736 world. *Trends Cogn Sci* 10:294–300.
- 13 737 Daniel R, Wagner G, Koch K, Reichenbach JR, Sauer H, Schlösser RGM (2011): Assessing the
- 14 738 neural basis of uncertainty in perceptual category learning through varying levels of
- 15 739 distortion. *J Cogn Neurosci* 23:1781–1793.
- 16 740 D’Ardenne K, McClure SM, Nystrom LE, Cohen JD (2008): BOLD responses reflecting
- 17 741 dopaminergic signals in the human ventral tegmental area. *Science* 319:1264–1267.
- 18 742 Daw ND, Niv Y, Dayan P (2005): Uncertainty-based competition between prefrontal and
- 19 743 dorsolateral striatal systems for behavioral control. *Nat Neurosci* 8:1704–1711.
- 20 744 Daw ND, Gershman SJ, Seymour B, Dayan P, Dolan RJ (2011): Model-Based Influences on
- 21 745 Humans’ Choices and Striatal Prediction Errors. *Neuron* 69:1204–1215.
- 22 746 Dayan P, Balleine BW (2002): Reward, motivation, and reinforcement learning. *Neuron*
- 23 747 36:285–298.
- 24 748 Delgado MR, Nystrom LE, Fissell C, Noll DC, Fiez JA (2000): Tracking the Hemodynamic
- 25 749 Responses to Reward and Punishment in the Striatum. *J Neurophysiol* 84:3072–
- 26 750 3077.
- 27 751 Delgado MR (2007): Reward-Related Responses in the Human Striatum. *Ann N Y Acad Sci*
- 28 752 1104:70–88.
- 29 753 Demos KE, Heatherton TF, Kelley WM (2012): Individual Differences in Nucleus Accumbens
- 30 754 Activity to Food and Sexual Images Predict Weight Gain and Sexual Behavior. *J*
- 31 755 *Neurosci* 32:5549–5552.
- 32 756 Diederer KJM, Ziauddeen H, Vestergaard MD, Spencer T, Schultz W, Fletcher PC (2017):
- 33 757 Dopamine Modulates Adaptive Prediction Error Coding in the Human Midbrain and
- 34 758 Striatum. *J Neurosci Off J Soc Neurosci* 37:1708–1720.
- 35 759 Diuk C, Tsai K, Wallis J, Botvinick M, Niv Y (2013): Hierarchical Learning Induces Two
- 36 760 Simultaneous, But Separable, Prediction Errors in Human Basal Ganglia. *J Neurosci*
- 37 761 33:5797–5805.
- 38 762 Doya K, Samejima K, Katagiri K, Kawato M (2002): Multiple model-based reinforcement
- 39 763 learning. *Neural Comput* 14:1347–1369.
- 40 764 Dreher J-C (2013): Neural coding of computational factors affecting decision making. *Prog*
- 41 765 *Brain Res* 202:289–320.
- 42 766 van Duijvenvoorde ACK, Op de Macks ZA, Overgaauw S, Gunther Moor B, Dahl RE, Crone EA
- 43 767 (2014): A cross-sectional and longitudinal analysis of reward-related brain activation:
- 44 768 Effects of age, pubertal stage, and reward sensitivity. *Brain Cogn* 89:3–14.
- 45 769 Dunne S, D’Souza A, O’Doherty JP (2016): The involvement of model-based but not model-
- 46 770 free learning signals during observational reward learning in the absence of choice. *J*
- 47 771 *Neurophysiol* 115:3195–3203.
- 48 772 Eickhoff SB, Etkin A (2016): Going Beyond Finding the “Lesion”: A Path for Maturation of
- 49 773 *Neuroimaging*. *Am J Psychiatry* 173:302–303.

- 1
2
3 774 Eickhoff SB, Laird AR, Fox PM, Lancaster JL, Fox PT (2017): Implementation errors in the
4 775 GingerALE Software: Description and recommendations. *Hum Brain Mapp* 38:7–11.
5 776 Eickhoff SB, Laird AR, Grefkes C, Wang LE, Zilles K, Fox PT (2009): Coordinate-based
6 777 activation likelihood estimation meta-analysis of neuroimaging data: a random-
7 778 effects approach based on empirical estimates of spatial uncertainty. *Hum Brain*
8 779 *Mapp* 30:2907–2926.
9
10 780 Elliot AJ (2006): The Hierarchical Model of Approach-Avoidance Motivation. *Motiv Emot*
11 781 30:111–116.
12 782 Elliott R, Friston KJ, Dolan RJ (2000): Dissociable neural responses in human reward systems.
13 783 *J Neurosci Off J Soc Neurosci* 20:6159–6165.
14 784 Elward RL, Vilberg KL, Rugg MD (2015): Motivated Memories: Effects of Reward and
15 785 Recollection in the Core Recollection Network and Beyond. *Cereb Cortex* 25:3159–
16 786 3166.
17
18 787 Ernst M, Nelson EE, McClure EB, Monk CS, Munson S, Eshel N, Zarah E, Leibenluft E,
19 788 Zametkin A, Towbin K, Blair J, Charney D, Pine DS (2004): Choice selection and
20 789 reward anticipation: an fMRI study. *Neuropsychologia* 42:1585–1597.
21 790 Ferdinand NK, Opitz B (2014): Different aspects of performance feedback engage different
22 791 brain areas: Disentangling valence and expectancy in feedback processing. *Sci Rep*
23 792 4:srep05986.
24
25 793 Fiorillo CD (2013): Two Dimensions of Value: Dopamine Neurons Represent Reward But Not
26 794 Aversiveness. *Science* 341:546–549.
27 795 Fiorillo CD, Tobler PN, Schultz W (2003): Discrete Coding of Reward Probability and
28 796 Uncertainty by Dopamine Neurons. *Science* 299:1898–1902.
29 797 Forster SE, Brown JW (2011): Medial prefrontal cortex predicts and evaluates the timing of
30 798 action outcomes. *NeuroImage* 55:253–265.
31
32 799 Fouragnan E, Chierchia G, Greiner S, Neveu R, Avesani P, Coricelli G (2013): Reputational
33 800 Priors Magnify Striatal Responses to Violations of Trust. *J Neurosci* 33:3602–3611.
34 801 Fouragnan E, Queirazza F, Retzler C, Mullinger K, Philiastides M (2017): Spatiotemporal
35 802 neural characterization of prediction error valence and surprise during reward
36 803 learning in humans. *Sci Rep*. <https://www.nature.com/srep/>.
37 804 Fouragnan E, Retzler C, Mullinger K, Philiastides MG (2015): Two spatiotemporally distinct
38 805 value systems shape reward-based learning in the human brain. *Nat Commun*
39 806 6:8107.
40
41 807 Frank MJ, Seeberger LC, O'reilly RC (2004): By carrot or by stick: cognitive reinforcement
42 808 learning in parkinsonism. *Science* 306:1940–1943.
43 809 Friston K (2009): The free-energy principle: a rough guide to the brain? *Trends Cogn Sci*
44 810 13:293–301.
45 811 Friston K, Kilner J, Harrison L (2006): A free energy principle for the brain. *J Physiol Paris*
46 812 100:70–87.
47
48 813 Fujiwara J, Tobler PN, Taira M, Iijima T, Tsutsui K-I (2009): Segregated and integrated coding
49 814 of reward and punishment in the cingulate cortex. *J Neurophysiol* 101:3284–3293.
50 815 Garrison J, Erdeniz B, Done J (2013): Prediction error in reinforcement learning: A meta-
51 816 analysis of neuroimaging studies. *Neurosci Biobehav Rev* 37:1297–1310.
52 817 Gherman S, Philiastides MG (2015): Neural representations of confidence emerge from the
53 818 process of decision formation during perceptual choices. *NeuroImage* 106:134–143.
54 819 Gherman S, Philiastides MG (2017): Human VMPFC encodes early signatures of confidence
55 820 in perceptual decisions. *bioRxiv*:224337.
56
57
58
59
60

- 1
- 2
- 3 821 Gläscher J, Daw N, Dayan P, O'Doherty JP (2010): States versus rewards: dissociable neural
- 4 822 prediction error signals underlying model-based and model-free reinforcement
- 5 823 learning. *Neuron* 66:585–595.
- 6 824 Gläscher J, Hampton AN, O'Doherty JP (2009): Determining a Role for Ventromedial
- 7 825 Prefrontal Cortex in Encoding Action-Based Value Signals During Reward-Related
- 8 826 Decision Making. *Cereb Cortex* 19:483–495.
- 9 827 Gray JA, McNaughton N (2003): *The Neuropsychology of Anxiety: An Enquiry Into the*
- 10 828 *Function of the Septo-hippocampal System*. Oxford University Press.
- 11 829 Guitart-Masip M, Huys QJM, Fuentemilla L, Dayan P, Duzel E, Dolan RJ (2012): Go and no-go
- 12 830 learning in reward and punishment: interactions between affect and effect.
- 13 831 *NeuroImage* 62:154–166.
- 14 832 Guo R, Böhmer W, Hebart M, Chien S, Sommer T, Obermayer K, Gläscher J (2016):
- 15 833 Interaction of Instrumental and Goal-Directed Learning Modulates Prediction Error
- 16 834 Representations in the Ventral Striatum. *J Neurosci* 36:12650–12660.
- 17 835 Haber SN, Kunishio K, Mizobuchi M, Lynd-Balta E (1995): The orbital and medial prefrontal
- 18 836 circuit through the primate basal ganglia. *J Neurosci Off J Soc Neurosci* 15:4851–
- 19 837 4867.
- 20 838 Hajcak G, Moser JS, Holroyd CB, Simons RF (2006): The feedback-related negativity reflects
- 21 839 the binary evaluation of good versus bad outcomes. *Biol Psychol* 71:148–154.
- 22 840 Hall G, Pearce JM (1979): Latent inhibition of a CS during CS-US pairings. *J Exp Psychol Anim*
- 23 841 *Behav Process* 5:31–42.
- 24 842 Hare TA, O'Doherty J, Camerer CF, Schultz W, Rangel A (2008): Dissociating the Role of the
- 25 843 Orbitofrontal Cortex and the Striatum in the Computation of Goal Values and
- 26 844 Prediction Errors. *J Neurosci* 28:5623–5630.
- 27 845 Haruno M, Kuroda T, Doya K, Toyama K, Kimura M, Samejima K, Imamizu H, Kawato M
- 28 846 (2004): A neural correlate of reward-based behavioral learning in caudate nucleus: a
- 29 847 functional magnetic resonance imaging study of a stochastic decision task. *J Neurosci*
- 30 848 *Off J Soc Neurosci* 24:1660–1665.
- 31 849 Häusler AN, Oroz Artigas S, Trautner P, Weber B (2016): Gain- and Loss-Related Brain
- 32 850 Activation Are Associated with Information Search Differences in Risky Gambles: An
- 33 851 fMRI and Eye-Tracking Study. *eNeuro* 3.
- 34 852 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5032244/>.
- 35 853 Hayden BY, Heilbronner SR, Pearson JM, Platt ML (2011): Surprise Signals in Anterior
- 36 854 Cingulate Cortex: Neuronal Encoding of Unsigned Reward Prediction Errors Driving
- 37 855 Adjustment in Behavior. *J Neurosci* 31:4178–4187.
- 38 856 Hester R, Barre N, Murphy K, Silk TJ, Mattingley JB (2008): Human Medial Frontal Cortex
- 39 857 Activity Predicts Learning from Errors. *Cereb Cortex* 18:1933–1940.
- 40 858 Hester R, Murphy K, Brown FL, Skilleter AJ (2010): Punishing an Error Improves Learning: The
- 41 859 Influence of Punishment Magnitude on Error-Related Neural Activity and Subsequent
- 42 860 Learning. *J Neurosci* 30:15600–15607.
- 43 861 Huettel SA, Stowe CJ, Gordon EM, Warner BT, Platt ML (2006): Neural signatures of
- 44 862 economic preferences for risk and ambiguity. *Neuron* 49:765–775.
- 45 863 Ide JS, Shenoy P, Yu AJ, Li CR (2013): Bayesian Prediction and Evaluation in the Anterior
- 46 864 Cingulate Cortex. *J Neurosci* 33:2039–2047.
- 47 865 Iglesias S, Mathys C, Brodersen KH, Kasper L, Piccirelli M, den Ouden HEM, Stephan KE
- 48 866 (2013): Hierarchical Prediction Errors in Midbrain and Basal Forebrain during Sensory
- 49 867 Learning. *Neuron* 80:519–530.
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- Ikemoto S (2007): Dopamine reward circuitry: Two projection systems from the ventral midbrain to the nucleus accumbens–olfactory tubercle complex. *Brain Res Rev* 56:27–78.
- Jenkinson M, Bannister P, Brady M, Smith S (2002): Improved optimization for the robust and accurate linear registration and motion correction of brain images. *NeuroImage* 17:825–841.
- Jensen J, Smith AJ, Willeit M, Crawley AP, Mikulis DJ, Vitcu I, Kapur S (2007): Separate brain regions code for salience vs. valence during reward prediction in humans. *Hum Brain Mapp* 28:294–302.
- Jocham G, Brodersen KH, Constantinescu AO, Kahn MC, Ianni AM, Walton ME, Rushworth MFS, Behrens TEJ (2016): Reward-Guided Learning with and without Causal Attribution. *Neuron* 90:177–190.
- Kaelbling LP, Littman ML, Moore AW (1996): Reinforcement Learning: A Survey. *arXiv:cs/9605103*. <http://arxiv.org/abs/cs/9605103>.
- Kahnt T (2017): A decade of decoding reward-related fMRI signals and where we go from here. *NeuroImage*. <http://www.sciencedirect.com/science/article/pii/S1053811917304688>.
- Kahnt T, Heinzle J, Park SQ, Haynes J-D (2010): The neural code of reward anticipation in human orbitofrontal cortex. *Proc Natl Acad Sci* 107:6010–6015.
- Kahnt T, Tobler PN (2013): Salience Signals in the Right Temporoparietal Junction Facilitate Value-Based Decisions. *J Neurosci* 33:863–869.
- Katahira K, Matsuda Y-T, Fujimura T, Ueno K, Asamizuya T, Suzuki C, Cheng K, Okanoya K, Okada M (2015): Neural basis of decision making guided by emotional outcomes. *J Neurophysiol* 113:3056–3068.
- Klein-Flügge MC, Hunt LT, Bach DR, Dolan RJ, Behrens TEJ (2011): Dissociable reward and timing signals in human midbrain and ventral striatum. *Neuron* 72:654–664.
- Knutson B, Adams CM, Fong GW, Hommer D (2001): Anticipation of increasing monetary reward selectively recruits nucleus accumbens. *J Neurosci Off J Soc Neurosci* 21:RC159.
- Knutson B, Westdorp A, Kaiser E, Hommer D (2000): FMRI visualization of brain activity during a monetary incentive delay task. *NeuroImage* 12:20–27.
- Koch K, Schachtzabel C, Wagner G, Reichenbach JR, Sauer H, Schlösser R (2008): The neural correlates of reward-related trial-and-error learning: An fMRI study with a probabilistic learning task. *Learn Mem* 15:728–732.
- Kolling N, Behrens TEJ, Mars RB, Rushworth MFS (2012): Neural Mechanisms of Foraging. *Science* 336:95–98.
- Kotz SA, Dengler R, Wittfoth M (2015): Valence-specific conflict moderation in the dorso-medial PFC and the caudate head in emotional speech. *Soc Cogn Affect Neurosci* 10:165–171.
- Kurniawan IT, Guitart-Masip M, Dayan P, Dolan RJ (2013): Effort and valuation in the brain: the effects of anticipation and execution. *J Neurosci Off J Soc Neurosci* 33:6160–6169.
- Laird AR, Fox PM, Price CJ, Glahn DC, Uecker AM, Lancaster JL, Turkeltaub PE, Kochunov P, Fox PT (2005): ALE meta-analysis: controlling the false discovery rate and performing statistical contrasts. *Hum Brain Mapp* 25:155–164.
- Lebreton M, Abitbol R, Daunizeau J, Pessiglione M (2015): Automatic integration of confidence in the brain valuation signal. *Nat Neurosci* 18:1159–1167.

- 1
- 2
- 3 915 Leknes S, Lee M, Berna C, Andersson J, Tracey I (2011): Relief as a Reward: Hedonic and
- 4 916 Neural Responses to Safety from Pain. *PLOS ONE* 6:e17870.
- 5 917 Leong YC, Radulescu A, Daniel R, DeWoskin V, Niv Y (2017): Dynamic Interaction between
- 6 918 Reinforcement Learning and Attention in Multidimensional Environments. *Neuron*
- 7 919 93:451–463.
- 8 920 Li J, Daw ND (2011): Signals in human striatum are appropriate for policy update rather than
- 9 921 value prediction. *J Neurosci Off J Soc Neurosci* 31:5504–5511.
- 10 922 Li J, Zhang Y (2006): Prediction error method-based second-order structural identification
- 11 923 algorithm in stochastic state space formulation. *Earthq Eng Struct Dyn* 35:761–779.
- 12 924 Lin A, Adolphs R, Rangel A (2012): Social and monetary reward learning engage overlapping
- 13 925 neural substrates. *Soc Cogn Affect Neurosci* 7:274–281.
- 14 926 Litt A, Plassmann H, Shiv B, Rangel A (2011): Dissociating valuation and saliency signals
- 15 927 during decision-making. *Cereb Cortex N Y N 1991* 21:95–102.
- 16 928 Liu X, Hairston J, Schrier M, Fan J (2011): Common and distinct networks underlying reward
- 17 929 valence and processing stages: a meta-analysis of functional neuroimaging studies.
- 18 930 *Neurosci Biobehav Rev* 35:1219–1236.
- 19 931 Losecaat Vermeer A, Boksem M, G Sanfey A (2014): Neural mechanisms underlying context-
- 20 932 dependent shifts in risk preferences. *NeuroImage* 103.
- 21 933 Luking KR, Luby JL, Barch DM (2014): Kids, candy, brain and behavior: Age differences in
- 22 934 responses to candy gains and losses. *Dev Cogn Neurosci* 9:82–92.
- 23 935 Manza P, Hu S, Ide JS, Farr OM, Zhang S, Leung H-C, Li CR (2016): The effects of
- 24 936 methylphenidate on cerebral responses to conflict anticipation and unsigned
- 25 937 prediction error in a stop-signal task. *J Psychopharmacol Oxf Engl* 30:283–293.
- 26 938 Marsh R, Hao X, Xu D, Wang Z, Duan Y, Liu J, Kangarlu A, Martinez D, Garcia F, Tau GZ, Yu S,
- 27 939 Packard MG, Peterson BS (2010): A virtual reality-based fMRI study of reward-based
- 28 940 spatial learning. *Neuropsychologia* 48:2912–2921.
- 29 941 Mathys CD, Lomakina EI, Daunizeau J, Iglesias S, Brodersen KH, Friston KJ, Stephan KE
- 30 942 (2014): Uncertainty in perception and the Hierarchical Gaussian Filter. *Front Hum*
- 31 943 *Neurosci* 8. <http://journal.frontiersin.org/article/10.3389/fnhum.2014.00825/full>.
- 32 944 Matsumoto M, Hikosaka O (2009): Two types of dopamine neuron distinctly convey positive
- 33 945 and negative motivational signals. *Nature* 459:837–841.
- 34 946 Mattfeld AT, Gluck MA, Stark CEL (2011): Functional specialization within the striatum along
- 35 947 both the dorsal/ventral and anterior/posterior axes during associative learning via
- 36 948 reward and punishment. *Learn Mem Cold Spring Harb N* 18:703–711.
- 37 949 Maunsell JHR (2004): Neuronal representations of cognitive state: reward or attention?
- 38 950 *Trends Cogn Sci* 8:261–265.
- 39 951 McClure SM, Berns GS, Montague PR (2003): Temporal prediction errors in a passive
- 40 952 learning task activate human striatum. *Neuron* 38:339–346.
- 41 953 Menon V, Levitin DJ (2005): The rewards of music listening: response and physiological
- 42 954 connectivity of the mesolimbic system. *NeuroImage* 28:175–184.
- 43 955 Metereau E, Dreher J-C (2013): Cerebral correlates of salient prediction error for different
- 44 956 rewards and punishments. *Cereb Cortex N Y N 1991* 23:477–487.
- 45 957 Metereau E, Dreher J-C (2015): The medial orbitofrontal cortex encodes a general unsigned
- 46 958 value signal during anticipation of both appetitive and aversive events. *Cortex J*
- 47 959 *Devoted Study Nerv Syst Behav* 63:42–54.
- 48 960 Meyniel F, Dehaene S (2017): Brain networks for confidence weighting and hierarchical
- 49 961 inference during probabilistic learning. *Proc Natl Acad Sci* 114:E3859–E3868.
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 962 Middlebrooks PG, Sommer MA (2012): Neuronal correlates of metacognition in primate
4 963 frontal cortex. *Neuron* 75:517–530.
- 5 964 Montague PR, Dayan P, Sejnowski TJ (1996): A framework for mesencephalic dopamine
6 965 systems based on predictive Hebbian learning. *J Neurosci Off J Soc Neurosci*
7 966 16:1936–1947.
- 8 967 Morrens J (2014): Dopamine neurons coding prediction errors in reward space, but not in
9 968 aversive space: a matter of location? *J Neurophysiol* 112:1021–1024.
- 10 969 Nassar MR, Rumsey KM, Wilson RC, Parikh K, Heasley B, Gold JI (2012): Rational regulation of
11 970 learning dynamics by pupil-linked arousal systems. *Nat Neurosci* 15:1040–1046.
- 12 971 Nieuwenhuis S, Slagter HA, von Geusau NJA, Heslenfeld DJ, Holroyd CB (2005): Knowing
13 972 good from bad: differential activation of human cortical areas by positive and
14 973 negative outcomes. *Eur J Neurosci* 21:3161–3168.
- 15 974 Niv Y, Daniel R, Geana A, Gershman SJ, Leong YC, Radulescu A, Wilson RC (2015):
16 975 Reinforcement Learning in Multidimensional Environments Relies on Attention
17 976 Mechanisms. *J Neurosci* 35:8145–8157.
- 18 977 Noonan MP, Mars RB, Rushworth MFS (2011): Distinct roles of three frontal cortical areas in
19 978 reward-guided behavior. *J Neurosci Off J Soc Neurosci* 31:14399–14412.
- 20 979 O'Doherty J, Kringelbach ML, Rolls ET, Hornak J, Andrews C (2001): Abstract reward and
21 980 punishment representations in the human orbitofrontal cortex. *Nat Neurosci* 4:95–
22 981 102.
- 23 982 O'Doherty J, Critchley H, Deichmann R, Dolan RJ (2003): Dissociating valence of outcome
24 983 from behavioral control in human orbital and ventral prefrontal cortices. *J Neurosci*
25 984 *Off J Soc Neurosci* 23:7931–7939.
- 26 985 O'Doherty J, Dayan P, Schultz J, Deichmann R, Friston K, Dolan RJ (2004): Dissociable Roles
27 986 of Ventral and Dorsal Striatum in Instrumental Conditioning. *Science* 304:452–454.
- 28 987 O'doherty JP, Hampton A, Kim H (2007): Model-Based fMRI and Its Application to Reward
29 988 Learning and Decision Making. *Ann N Y Acad Sci* 1104:35–53.
- 30 989 Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011): FieldTrip: Open Source Software for
31 990 Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. Research
32 991 article. *Computational Intelligence and Neuroscience*.
33 992 <https://www.hindawi.com/journals/cin/2011/156869/>.
- 34 993 O'Reilly JX, Schüffegen U, Cuell SF, Behrens TEJ, Mars RB, Rushworth MFS (2013):
35 994 Dissociable effects of surprise and model update in parietal and anterior cingulate
36 995 cortex. *Proc Natl Acad Sci* 110:E3660–E3669.
- 37 996 den Ouden HEM, Kok P, de Lange FP (2012): How Prediction Errors Shape Perception,
38 997 Attention, and Motivation. *Front Psychol* 3.
39 998 <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3518876/>.
- 40 999 Pagnoni G, Zink CF, Montague PR, Berns GS (2002): Activity in human ventral striatum
41 1000 locked to errors of reward prediction. *Nat Neurosci* 5:97–98.
- 42 1001 Paschke LM, Walter H, Steimke R, Ludwig VU, Gaschler R, Schubert T, Stelzel C (2015):
43 1002 Motivation by potential gains and losses affects control processes via different
44 1003 mechanisms in the attentional network. *NeuroImage* 111:549–561.
- 45 1004 Pearce JM, Hall G (1980): A model for Pavlovian learning: Variations in the effectiveness of
46 1005 conditioned but not of unconditioned stimuli. *Psychol Rev* 87:532–552.
- 47 1006 Pessiglione M, Petrovic P, Daunizeau J, Palminteri S, Dolan RJ, Frith CD (2008): Subliminal
48 1007 instrumental conditioning demonstrated in the human brain. *Neuron* 59:561–567.

- 1008 Pessiglione M, Seymour B, Flandin G, Dolan RJ, Frith CD (2006): Dopamine-dependent
 1009 prediction errors underpin reward-seeking behaviour in humans. *Nature* 442:1042–
 1010 1045.
- 1011 Philiastides MG, Biele G, Heekeren HR (2010a): A mechanistic account of value computation
 1012 in the human brain. *Proc Natl Acad Sci U S A* 107:9430–9435.
- 1013 Philiastides MG, Biele G, Vavatzanidis N, Kazzner P, Heekeren HR (2010b): Temporal dynamics
 1014 of prediction error processing during reward-based decision making. *NeuroImage*
 1015 53:221–232.
- 1016 Philiastides MG, Heekeren HR, Sajda P (2014): Human Scalp Potentials Reflect a Mixture of
 1017 Decision-Related Signals during Perceptual Choices. *J Neurosci* 34:16877–16889.
- 1018 Pisauro A, Fouragnan E, Retzler C, Philiastides M (2017): Neural correlates of evidence
 1019 accumulation during value-based decisions revealed via simultaneous EEG-fMRI. *Nat*
 1020 *Commun.* <http://eprints.hud.ac.uk/31834/>.
- 1021 Poudel GR, Innes CRH, Jones RD (2013): Distinct neural correlates of time-on-task and
 1022 transient errors during a visuomotor tracking task after sleep restriction.
 1023 *NeuroImage* 77:105–113.
- 1024 Preuschoff K, Quartz SR, Bossaerts P (2008): Human Insula Activation Reflects Risk
 1025 Prediction Errors As Well As Risk. *J Neurosci* 28:2745–2752.
- 1026 Queirazza F, Fouragnan E, Steele JD, Cavanagh J, Philiastides M (2017): Dorsomedial
 1027 prefrontal cortex activity during reinforcement learning discriminates response to
 1028 Cognitive Behavioural Therapy in depression. *bioRxiv*:224410.
- 1029 R Garrison J, Done J, S Simons J (2017): Interpretation of published meta-analytical studies
 1030 affected by implementation errors in the GingerALE software.
- 1031 Rescorla R, Wagner A (1972): A theory of Pavlovian conditioning: Variations in the
 1032 effectiveness of reinforcement and nonreinforcement. In: Black, A, Prokasy, W,
 1033 editors. *Classical Conditioning II: Current Research and Theory*. Appleton-Century-
 1034 Crofts. pp 64–99.
- 1035 Ribas-Fernandes JF, Solway A, Diuk C, McGuire JT, Barto AG, Niv Y, Botvinick MM (2011): A
 1036 neural signature of hierarchical reinforcement learning. *Neuron* 71:370–379.
- 1037 Rodriguez PF (2009): Stimulus-outcome learnability differentially activates anterior cingulate
 1038 and hippocampus at feedback processing. *Learn Mem Cold Spring Harb N* 16:324–
 1039 331.
- 1040 Roesch MR, Olson CR (2004): Neuronal Activity Related to Reward Value and Motivation in
 1041 Primate Frontal Cortex. *Science* 304:307–310.
- 1042 Rohe T, Noppeney U (2015): Cortical Hierarchies Perform Bayesian Causal Inference in
 1043 Multisensory Perception. *PLOS Biol* 13:e1002073.
- 1044 Rohe T, Weber B, Fliessbach K (2012): Dissociation of BOLD responses to reward prediction
 1045 errors and reward receipt by a model comparison. *Eur J Neurosci* 36:2376–2382.
- 1046 Rolls ET, Grabenhorst F, Parris BA (2008): Warm pleasant feelings in the brain. *NeuroImage*
 1047 41:1504–1513.
- 1048 Rudolf S, Preuschoff K, Weber B (2012): Neural Correlates of Anticipation Risk Reflect Risk
 1049 Preferences. *J Neurosci* 32:16683–16692.
- 1050 Samson RD, Frank MJ, Fellous J-M (2010): Computational models of reinforcement learning:
 1051 the role of dopamine as a reward signal. *Cogn Neurodyn* 4:91–105.
- 1052 Sarinopoulos I, Grupe DW, Mackiewicz KL, Herrington JD, Lor M, Steege EE, Nitschke JB
 1053 (2010): Uncertainty during anticipation modulates neural responses to aversion in
 1054 human insula and amygdala. *Cereb Cortex N Y N 1991* 20:929–940.

- 1055 Schlagenhauf F, Rapp MA, Huys QJM, Beck A, Wüstenberg T, Deserno L, Buchholz H-G,
 1056 Kalbitzer J, Buchert R, Bauer M, Kienast T, Cumming P, Plotkin M, Kumakura Y, Grace
 1057 AA, Dolan RJ, Heinz A (2013): Ventral striatal prediction error signaling is associated
 1058 with dopamine synthesis capacity and fluid intelligence. *Hum Brain Mapp* 34:1490–
 1059 1499.
- 1060 Scholl J, Kolling N, Nelissen N, Wittmann MK, Harmer CJ, Rushworth MFS (2015): The Good,
 1061 the Bad, and the Irrelevant: Neural Mechanisms of Learning Real and Hypothetical
 1062 Rewards and Effort. *J Neurosci Off J Soc Neurosci* 35:11233–11251.
- 1063 Schonberg T, O'Doherty JP, Joel D, Inzelberg R, Segev Y, Daw ND (2010): Selective
 1064 impairment of prediction error signaling in human dorsolateral but not ventral
 1065 striatum in Parkinson's disease patients: evidence from a model-based fMRI study.
 1066 *NeuroImage* 49:772–781.
- 1067 Schultz W, Apicella P, Ljungberg T (1993): Responses of monkey dopamine neurons to
 1068 reward and conditioned stimuli during successive steps of learning a delayed
 1069 response task. *J Neurosci Off J Soc Neurosci* 13:900–913.
- 1070 Schultz W, Apicella P, Scarnati E, Ljungberg T (1992): Neuronal activity in monkey ventral
 1071 striatum related to the expectation of reward. *J Neurosci Off J Soc Neurosci*
 1072 12:4595–4610.
- 1073 Schultz W (1998): Predictive Reward Signal of Dopamine Neurons. *J Neurophysiol* 80:1–27.
- 1074 Schultz W (2016a): Dopamine reward prediction-error signalling: a two-component
 1075 response. *Nat Rev Neurosci* 17:183–195.
- 1076 Schultz W (2016b): Dopamine reward prediction error coding. *Dialogues Clin Neurosci*
 1077 18:23–32.
- 1078 Schultz W, Dayan P, Montague PR (1997): A Neural Substrate of Prediction and Reward.
 1079 *Science* 275:1593–1599.
- 1080 Schwartenbeck P, FitzGerald THB, Dolan R (2016): Neural signals encoding shifts in beliefs.
 1081 *NeuroImage* 125:578–586.
- 1082 Scimeca JM, Katzman PL, Badre D (2016): Striatal prediction errors support dynamic control
 1083 of declarative memory decisions. *Nat Commun* 7:ncomms13061.
- 1084 Serences JT (2008): Value-based modulations in human visual cortex. *Neuron* 60:1169–
 1085 1181.
- 1086 Sescousse G, Caldú X, Segura B, Dreher J-C (2013): Processing of primary and secondary
 1087 rewards: A quantitative meta-analysis and review of human functional neuroimaging
 1088 studies. *Neurosci Biobehav Rev* 37:681–696.
- 1089 Seymour B, Daw N, Dayan P, Singer T, Dolan R (2007): Differential Encoding of Losses and
 1090 Gains in the Human Striatum. *J Neurosci Off J Soc Neurosci* 27:4826–4831.
- 1091 Silvetti M, Verguts T (2012): Reinforcement Learning, High-Level Cognition, and the Human
 1092 Brain. [http://www.intechopen.com/books/neuroimaging-cognitive-and-clinical-](http://www.intechopen.com/books/neuroimaging-cognitive-and-clinical-neuroscience/reinforcement-learning-high-level-cognition-and-the-human-brain)
 1093 [neuroscience/reinforcement-learning-high-level-cognition-and-the-human-brain.](http://www.intechopen.com/books/neuroimaging-cognitive-and-clinical-neuroscience/reinforcement-learning-high-level-cognition-and-the-human-brain)
- 1094 So N, Stuphorn V (2012): Supplementary Eye Field Encodes Reward Prediction Error. *J*
 1095 *Neurosci* 32:2950–2963.
- 1096 Späti J, Chumbley J, Brakowski J, Dörig N, Grosse Holtforth M, Seifritz E, Spinelli S (2014):
 1097 Functional lateralization of the anterior insula during feedback processing. *Hum*
 1098 *Brain Mapp* 35:4428–4439.
- 1099 Spicer J, Galvan A, Hare TA, Voss H, Glover G, Casey B (2007): Sensitivity of the Nucleus
 1100 Accumbens to Violations in Expectation of Reward. *NeuroImage* 34:455–461.

- 1101 Spoormaker VI, Andrade KC, Schröter MS, Sturm A, Goya-Maldonado R, Sämann PG, Czisch
- 1102 M (2011): The neural correlates of negative prediction error signaling in human fear
- 1103 conditioning. *NeuroImage* 54:2250–2256.
- 1104 Sutton R (1998): Reinforcement Learning: An Introduction. Cambridge, Mass: MIT Press.
- 1105 Takemura H, Samejima K, Vogels R, Sakagami M, Okuda J (2011): Stimulus-Dependent
- 1106 Adjustment of Reward Prediction Error in the Midbrain. *PLOS ONE* 6:e28337.
- 1107 Tanaka SC, Doya K, Okada G, Ueda K, Okamoto Y, Yamawaki S (2004): Prediction of
- 1108 immediate and future rewards differentially recruits cortico-basal ganglia loops. *Nat*
- 1109 *Neurosci* 7:nn1279.
- 1110 Tanaka SC, Samejima K, Okada G, Ueda K, Okamoto Y, Yamawaki S, Doya K (2006): Brain
- 1111 mechanism of reward prediction under predictable and unpredictable environmental
- 1112 dynamics. *Neural Netw Off J Int Neural Netw Soc* 19:1233–1241.
- 1113 Tobia MJ, Gläscher J, Sommer T (2016): Context-specific behavioral surprise is differentially
- 1114 correlated with activity in anterior and posterior brain systems. *Neuroreport* 27:677–
- 1115 682.
- 1116 Turkeltaub PE, Eden GF, Jones KM, Zeffiro TA (2002): Meta-analysis of the functional
- 1117 neuroanatomy of single-word reading: method and validation. *NeuroImage* 16:765–
- 1118 780.
- 1119 Ullsperger M, Cramon DY von (2003): Error Monitoring Using External Feedback: Specific
- 1120 Roles of the Habenular Complex, the Reward System, and the Cingulate Motor Area
- 1121 Revealed by Functional Magnetic Resonance Imaging. *J Neurosci* 23:4308–4314.
- 1122 Valentin VV, O'Doherty JP (2009): Overlapping prediction errors in dorsal striatum during
- 1123 instrumental learning with juice and money reward in the human brain. *J*
- 1124 *Neurophysiol* 102:3384–3391.
- 1125 Watanabe N, Sakagami M, Haruno M (2013): Reward Prediction Error Signal Enhanced by
- 1126 Striatum–Amygdala Interaction Explains the Acceleration of Probabilistic Reward
- 1127 Learning by Emotion. *J Neurosci* 33:4487–4493.
- 1128 Wittmann MK, Kolling N, Akaishi R, Chau BKH, Brown JW, Nelissen N, Rushworth MFS
- 1129 (2016): Predictive decision making driven by multiple time-linked reward
- 1130 representations in the anterior cingulate cortex. *Nat Commun* 7:12327.
- 1131 Wunderlich K, Rangel A, O'Doherty JP (2009): Neural computations underlying action-based
- 1132 decision making in the human brain. *Proc Natl Acad Sci* 106:17199–17204.
- 1133 Wunderlich K, Symmonds M, Bossaerts P, Dolan RJ (2011): Hedging your bets by learning
- 1134 reward correlations in the human brain. *Neuron* 71:1141–1152.
- 1135 Yacubian J, Gläscher J, Schroeder K, Sommer T, Braus DF, Büchel C (2006): Dissociable
- 1136 systems for gain- and loss-related value predictions and errors of prediction in the
- 1137 human brain. *J Neurosci Off J Soc Neurosci* 26:9530–9537.
- 1138 Yeung N, Sanfey AG (2004): Independent Coding of Reward Magnitude and Valence in the
- 1139 Human Brain. *J Neurosci* 24:6258–6264.
- 1140 Yu AJ, Dayan P (2005): Uncertainty, Neuromodulation, and Attention. *Neuron* 46:681–692.
- 1141 Zalla T, Koechlin E, Pietrini P, Basso G, Aquino P, Sirigu A, Grafman J (2000): Differential
- 1142 amygdala responses to winning and losing: a functional magnetic resonance imaging
- 1143 study in humans. *Eur J Neurosci* 12:1764–1770.
- 1144 Zhang S, Mano H, Ganesh G, Robbins T, Seymour B (2016): Dissociable Learning Processes
- 1145 Underlie Human Pain Conditioning. *Curr Biol* 26:52–58.
- 1146 Zink CF, Pagnoni G, Martin-Skurski ME, Chappelw JC, Berns GS (2004): Human striatal
- 1147 responses to monetary reward depend on saliency. *Neuron* 42:509–517.

Table 1. Categorisation of fMRI studies into the three RPE components (valence, surprise, signed RPE) and broken down by the relevant fMRI contrast/regressor.

Statistical comparisons	Number	Total	Reference
Valence Pattern A i (NEG>POS)		32	[de Bruijn et al., 2009; Daniel et al., 2011; Demos et al., 2012; van Duijvenvoorde et al., 2014; Elward et al., 2015; Ferdinand and Opitz, 2014; Fouragnan et al., 2015; Gläscher et al., 2009; Haruno et al., 2004; Häusler et al., 2016; Jocham et al., 2016; Kahnt et al., 2010; Katahira et al., 2015; Klein-Flügge et al., 2011; Klein-Flügge et al., 2011; Knutson et al., 2000; Knutson et al., 2001; Koch et al., 2008; Leknes et al., 2011; Losecaat Vermeer et al., 2014; Marsh et al., 2010; Mattfeld et al., 2011; Noonan et al., 2011; O'Doherty et al., 2001; O'Doherty et al., 2003; Rodriguez, 2009; Rolls et al., 2008; Scholl et al., 2015; Seymour et al., 2007; Spicer et al., 2007; Spoomaker et al., 2011; Ullsperger and Cramon, 2003; Yacubian et al., 2006]
Negative > Positive	19		
Negative > No outcomes	9		
Negative correlation with a regressor defining valence RPE (with a binary modulation whereby positive RPE = 1, and negative RPE = -1)	4		
Valence Pattern A ii (POS>NEG)		33	[Amiez et al., 2012; Aron et al., 2004; Bickel et al., 2009; de Bruijn et al., 2009; Canessa et al., 2013; Daniel et al., 2011; van Duijvenvoorde et al., 2014; Elliott et al., 2000; Ernst et al., 2004; Forster and Brown, 2011; Fouragnan et al., 2015; Fujiwara et al., 2009; Häusler et al., 2016; Hester et al., 2008; Hester et al., 2010; Jocham et al., 2016; Katahira et al., 2015; Knutson et al., 2000; Knutson et al., 2001; Knutson et al., 2001; Kumiawan et al., 2013; Losecaat Vermeer et al., 2014; Luking et al., 2014; Paschke et al., 2015; Sarinopoulos et al., 2010; Scholl et al., 2015; Schonberg et al., 2010; Seymour et al., 2007; Späti et al., 2014; Spoomaker et al., 2011; Ullsperger and Cramon, 2003]
Positive > Negative	18		
Positive > No outcomes	9		
Positive correlation with a regressor defining valence RPE (with a binary modulation whereby positive RPE = 1, and negative RPE = -1)	6		
Surprise Pattern B		41	[Allen et al., 2016; Amado et al., 2016; Amiez et al., 2012; Boll et al., 2013; Browning et al., 2010; Chumbley et al., 2014; Daw et al., 2011; Dreher, 2013; Ferdinand and Opitz, 2014; Forster and Brown, 2011; Fouragnan et al., 2015; Fouragnan et al., 2017; Fujiwara et al., 2009; Ide et al., 2013; Iglesias et al., 2013; Jensen et al., 2007; Knutson et al., 2001; Kotz et al., 2015; Leong et al., 2017; Losecaat Vermeer et al., 2014; Manza et al., 2016; McClure et al., 2003; Metereau and Dreher, 2013; Metereau and Dreher, 2015; Meyniel and Dehaene, 2017; Nieuwenhuis et al., 2005; O'Reilly et al., 2013; den Ouden et al., 2012; Poudel et al., 2013; Rodriguez, 2009; Rohe et al., 2012; Rohe and Noppeney, 2015; Rohe and Noppeney, 2015; Rolls et al., 2008; Schwartenbeck et al., 2016; Silvetti and Verguts, 2012; Tobia et al., 2016; Watanabe et al., 2013; Wunderlich et al., 2009; Wunderlich et al., 2011; Yacubian et al., 2006; Zalla et al., 2000; Zhang et al., 2016]
Unsigned RPE ("RL surprise")	12		
Unsigned Bayesian RPE ("Volatility", "Bayesian surprise")	13		
Positive and Negative outcomes > No or low outcomes	9		
"Associability" term of the Pearce et Hall model	2		
Parametric changes in magnitude of surprising positive RPE (unsigned)	3		
Parametric changes in magnitude of surprising negative RPE (unsigned)	2		
Signed RPE Pattern C		38	[Abler et al., 2006; Behrens et al., 2007; van den Bos et al., 2012; Cohen and Ranganath, 2007; Daw et al., 2011; Delgado et al., 2000; Delgado, 2007; Diederer et al., 2017; Diuk et al., 2013; Dunne et al., 2016; Gläscher et al., 2010; Guo et al., 2016; Hare et al., 2008; Ide et al., 2013; Katahira et al., 2015; Leong et al., 2017; Li and Zhang, 2006; Lin et al., 2012; Mattfeld et al., 2011; McClure et al., 2003; Metereau and Dreher, 2013; Metereau and Dreher, 2015; O'Doherty et al., 2003; Pessiglione et al., 2006; Pessiglione et al., 2008; Ribas-Fernandes et al., 2011; Rolls et al., 2008; Schlagenhauf et al., 2013; Schonberg et al., 2010; Scimeca et al., 2016; Seymour et al., 2007; Takemura et al., 2011; Tanaka et al., 2004; Tanaka et al., 2006; Valentin and O'Doherty, 2009; Watanabe et al., 2013; Wunderlich et al., 2011]
Signed RPE (from model-free RL models)	16		
Signed RPE (from model-based RL models)	8		
Signed Bayesian RPE	10		
High positive RPEs > low positive RPEs > low negative RPEs > high negative RPEs	4		

Table 2. ALE cluster results for the valence analysis: Pattern A (i) and (ii) (FDR-ID $P < 0.05$, with a minimum volume cluster size of 50 mm^3).

Region	R/L	x	y	z	Cluster size	ALE score
Pattern A (i) NEG > POS						
Dorsomedial cingulate cortex (dMCC)	R	2	24	36	12712	0.051
Anterior Insula (aINS)	R	32	24	-2	6120	0.062
-	L	-32	22	-4	4880	0.056
Pallidum	R	12	8	4	3360	0.04
-	L	-14	6	2	2520	0.029
Middle Frontal Gyrus	R	38	4	32	3152	0.029
-	R	30	10	56	488	0.021
-	L	-28	12	60	104	0.019
Inferior Parietal Lobule (IPL)	R	40	-48	42	2416	0.039
-	L	-38	-48	42	2216	0.043
Middle Temporal Gyrus (MTG)	R	60	-28	-6	1192	0.031
Amygdala	R	18	-6	-12	704	0.024
Thalamus	L	-12	-12	10	624	0.025
-	L	-6	-26	8	280	0.023
Habenula	R	2	-20	-18	312	0.022
Dorsolateral Prefrontal Cortex (dlPFC)	L	-44	28	32	360	0.020
-	R	40	34	30	344	0.020
Fusiform Area	L	-40	-62	-10	272	0.023
Precentral Cortex	L	-52	0	34	256	0.021
Dorsomedial Orbitofrontal Cortex (dmOFC)	R	38	58	-2	192	0.020
Dorsomedial Prefrontal Cortex (dmPFC)	R	20	50	4	120	0.018
Superior Temporal Sulcus	R	58	-42	22	120	0.017
Pattern A (ii) (POS > NEG)						
Ventral striatum (vSTR)	L	-12	8	-4	4880	0.052
-	R	8	8	-2	2880	0.038
Ventromedial Prefrontal Cortex (vmPFC)	L	-2	42	0	3416	0.037
Posterior Cingulate Cortex (PCC)	L	0	-32	36	240	0.016
-	L	0	-36	26	88	0.014
Ventrolateral OFC (vlOFC)	R	32	44	-10	144	0.015
Dorsomedial Prefrontal Cortex (dmPFC)	L	-6	-56	14	96	0.016
Medial Prefrontal Cortex (mPFC)	L	-2	46	20	88	0.014

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. ALE clusters results for the surprise analysis (FDR-ID $P < 0.05$, with a minimum volume cluster size of 50 mm^3).

Region	R/L	x	y	z	Cluster size	ALE score
Anterior mid-cingulate Cortex (aMCC)	R	4	24	34	4072	0.029
Anterior Insula (aINS)	R	32	24	-4	2496	0.050
-	L	-32	20	-4	1544	0.038
Inferior Parietal Lobule (IPL)	R	40	-46	42	1672	0.033
-	L	-40	-48	42	568	0.025
Dorsal Striatum (dSTR)	R	12	8	4	1400	0.034
-	L	-14	10	2	1216	0.021
Middle Temporal Gyrus (MTG)	R	60	-28	-8	648	0.022
Lateral Inferior Frontal Cortex	R	52	10	18	488	0.025
Lateral Central Frontal Gyrus	L	-44	26	30	392	0.019
Precentral Gyrus	R	48	12	34	360	0.019
-	L	-52	0	34	224	0.020
Midbrain	R	2	-20	-18	304	0.021
Dorsal mid-cingulate cortex (dMCC)	R	12	14	42	224	0.019
Hippocampus	R	20	-6	-10	160	0.018
Fusiform Gyrus	L	-40	-60	-10	112	0.017
Mid Occipital Pole	L	-16	-90	-6	112	0.016
Superior Temporal Sulcus	R	60	-40	20	64	0.015

Table 4. ALE cluster results for the conjunction analysis of valence and surprise (FDR-ID $p < 0.05$, with a minimum volume cluster size of 50 mm^3).

Region	R/L	x	y	z	Cluster size	ALE score
Striatum (STR)	R	12	6	4	1082	0.031
-	L	-12	12	4	376	0.021
Anterior Insula (aINS)	L	-32	20	-6	453	0.018
Anterior Mid-cingulate cortex (aMCC)	R	3	22	37	221	0.014
Inferior Parietal Lobule	L	40	-46	42	327	0.014

Table 5. ALE cluster results for the contrast analyses of valence and surprise (FDR-pN $p < 0.05$, with a minimum volume cluster size of 50 mm^3).

Region	R/L	x	y	z	Cluster size	ALE score
Valence vs. Surprise						
Ventral Striatum (vSTR)	L	-10	8	-10	1096	3.29
ventromedial prefrontal cortex (vmPFC)	L	-2	44	0	256	3.29
Positive vs. Surprise						
Ventral Striatum (vSTR)	L	-12	-8	-8	1872	3.29
ventromedial prefrontal cortex (vmPFC)	R	0	46	0	512	3.29
Ventral Striatum (vSTR)	R	8	8	-6	168	3.29
Negative vs. Surprise						
Middle Insula (mINS)	R	40	10	2	544	3.29
Mid Cingulate Cortex (MCC)	R	6	20	42	144	3.29
Surprise vs. Valence						
Anterior Insula (aINS)	R	32	24	-4	1224	3.29
Anterior Insula (aINS)	L	-32	20	-2	112	3.29
Ventral Tegmental Area (VTA)	L	-6	-16	-10	96	3.29
Ventral Tegmental Area (VTA)	R	2	-20	-16	72	3.29
Occipital Lobe	R	24	-80	-6	72	3.29
Surprise vs. Positive						
Anterior Insula (aINS)	R	32	22	-2	1648	3.29
Middle Temporal Gyrus (MTG)	R	40	-46	42	1184	3.29
Anterior Insula (aINS)	L	-32	22	-2	1016	3.29
Inferior Frontal Gyrus	R	52	10	18	184	3.29
Supplementary Motor Area (SMA)	L	-2	12	52	160	3.29
Surprise vs. Negative						
Angular Gyrus	R	40	-46	40	248	3.29
Anterior Insula (aINS)	R	32	28	-6	80	3.29
Dorsal Striatum (dSTR)	R	12	10	2	56	3.29

1
2
3 1167 **Table 6.** ALE clusters results for the signed RPE studies (FDR-ID $p < 0.05$, with a minimum
4
5 1168 volume cluster size of 50 mm^3).

Region	R/L	x	y	z	Cluster size	ALE score
Striatum (STR) (encompasses left and right hemispheres)	R	12	10	-4	10888	0.053
Putamen	R	30	-6	8	688	0.024
Anterior Mid-cingulate Cortex (aMCC)	R	6	26	46	160	0.018
-	L	-2	14	40	120	0.016
Anterior Cingulate Cortex (ACC)	R	4	36	20	112	0.017
Ventromedial prefrontal (vmPFC)	L	0	34	0	64	0.015
Lateral Inferior Frontal Gyrus (LIIFC)	L	-46	4	24	64	0.016

17 1169
18
19 1170 **Table 7.** ALE cluster results for the contrast analyses of signed RPE and valence as well as
20
21 1171 signed RPE and surprise (FDR-pN $p < 0.05$, with a minimum volume cluster size of 50 mm^3).

Region	R/L	x	y	z	Cluster size	ALE score
Positive – Signed RPE						
Ventromedial Prefrontal Cortex (vmPFC)	R	2	44	-15	160	3.29
Signed RPE - Positive						
No significant						
Negative – Signed RPE						
Middle Insula (mINS)	R	40	12	0	528	3.29
Dorsal Middle Cingulate Cortex (dMCC)	R	6	22	36	208	3.29
Middle Insula (mINS)	L	-38	18	-4	184	3.29
Habenula	L	-2	-26	8	168	2.58
Thalamus	R	8	-10	5	96	2.58
Signed RPE - Negative						
Ventral Striatum (vSTR)	R	10	10	-6	2208	3.29
Valence – Signed RPE						
Ventromedial Prefrontal Cortex (vmPFC)	R	2	44	-12	760	3.29
Middle Insula (mINS)	R	40	12	2	568	2.58
Dorsal Middle Cingulate Cortex (dMCC)	R	6	24	38	480	2.58
Signed RPE - Valence						
Ventral Striatum (vSTR)	R	12	16	-2	184	3.29
Surprise – Signed RPE						
Anterior Insula (aINS)	L	-34	22	0	704	3.29
Anterior Midcingulate Cortex (aMCC)	R	0	14	52	136	3.29
Pre supplementary motor area (preSMA)	R	0	14	52	136	3.29
Anterior Insula (aINS)	R	38	18	-2	88	3.29

Signed RPE - Surprise						
Ventral Striatum (vSTR)	L	-10	8	-10	904	3.29
Ventral Striatum (vSTR)	R	12	14	-3	192	3.29
Ventral Striatum (vSTR)	R	4	6	-6	72	3.29

1172

For Peer Review

Figure Legends

Figure 1. Hypothesized profiles for BOLD responses as function of the three RPE components. Pattern A (i and ii) describe the two categorical valence responses (orange and blue colours indicate (i) responses being greater for negative compared to positive outcomes [NEG > POS] and (ii) responses being greater for positive compared to negative outcomes [POS > NEG]). Pattern B captures surprise effects with greater responses to higher outcome deviations from expectations, independent of the sign (valence) of the RPE. Pattern C shows a monotonically increasing response profile consistent with a signed RPE representation.

Figure 2. Results of whole-brain ALE analysis along the valence component. Overlays of brain areas activated by correlations with NEG > POS (blue) and POS > NEG (orange) (Pattern A (i) and (ii), respectively; Fig. 1) (P-values corrected with FDR-ID [FID] and FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³). Representative slices are shown with MNI coordinates given below each image.

Figure 3. Results of the whole brain ALE analysis for the surprise component of RPE (pattern B, Figure 1). Overlay of brain areas activated by all analyses representing direct or indirect measures of the surprise component of RPE (P-values corrected with FDR-ID [FID] and FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³). Representative slices are shown with MNI coordinates given below each image.

Figure 4. Results of the ALE conjunction analysis between valence and surprise (purple). The regions identified earlier with separate ALE analyses along the valence (NEG > POS: blue, POS > NEG: orange) and surprise (green) components are shown for comparison purposes. P-values were corrected with FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³ for the initial maps. Representative slices are shown with MNI coordinates given bellow each image.

Figure 5. Results of the ALE contrast analyses for [valence – surprise] (left panel) and [surprise – valence]. P-values were corrected with FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³ for the initial maps. Representative slices are shown with MNI coordinates given bellow each image.

Figure 6. Results of whole brain ALE analysis for signed RPE. Overlay of brain areas activated by positive correlation with signed RPE (P-values corrected with FDR-ID [FID] and FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³). Representative slices are shown with MNI coordinates given bellow each image.

Figure 7. Results of the ALE conjunction analysis for all components of RPE. Overlay of brain areas individually activated by (1) valence (orange), (2) surprise (green), and (3) signed RPE (red), with P-values corrected with FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³ for the initial maps. Importantly, the overlap between the three analyses, shown in white, also corresponds to the only cluster found for the ALE conjunction analysis between valence/surprise vs. signed RPE. MNI coordinates are given below each image.

Figure 8. Results of the ALE contrast analyses for [signed RPE – positive valence] (left panel), [signed RPE – negative valence] (middle panel) and [signed RPE – (positive + negative valence)] (right panel). P-values were corrected with FDR-pN [FRN] < 0.05 and a minimum cluster volume of 50 mm³ for the initial maps. Representative slices are shown with MNI coordinates given bellow each image.




fig.1

91x81mm (300 x 300 DPI)






fig.2

378x339mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




fig.4

203x100mm (300 x 300 DPI)

Conjunction Surprise & Valence




fig.4

285x172mm (300 x 300 DPI)




fig.5

252x115mm (300 x 300 DPI)




fig.6

207x98mm (300 x 300 DPI)

Conjunction Surprise & Valence & Signed PE




fig.7

164x66mm (300 x 300 DPI)




fig.8

366x400mm (300 x 300 DPI)